



HAL
open science

Multivalent Glycomimetics with Affinity and Selectivity toward Fucose-Binding Receptors from Emerging Pathogens

David Goyard, Veronica Baldoneschi, Annabelle Varrot, Michele Fiore, Anne Imberty, Barbara Richichi, Olivier Renaudet, Cristina Nativi

► **To cite this version:**

David Goyard, Veronica Baldoneschi, Annabelle Varrot, Michele Fiore, Anne Imberty, et al.. Multivalent Glycomimetics with Affinity and Selectivity toward Fucose-Binding Receptors from Emerging Pathogens. *Bioconjugate Chemistry*, 2018, 29 (1), pp.83-88. 10.1021/acs.bioconjchem.7b00616 . hal-02356130

HAL Id: hal-02356130

<https://hal.univ-grenoble-alpes.fr/hal-02356130v1>

Submitted on 18 Jan 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Multivalent Glycomimetics with Affinity and Selectivity Towards Fucose-binding Receptors from Emerging Pathogens

D. Goyard,[†] V. Baldoneschi,[‡] A. Varrot,[‡] M. Fiore,[‡] A. Imberty,^{‡*} B. Richichi,[‡] O. Renaudet^{†*} and
C. Nativi^{‡*}

[†] DCM, UMR5250, CNRS, University Grenoble Alpes, BP 53, 38041 Grenoble, France

[‡] Department of Chemistry Ugo Schiff, University of Florence, via della Lastruccia, 13 – 50019 Sesto F.no
(FI) Italy

[‡] CERMAV, UPR5301, CNRS University Grenoble Alpes BP 53, 38041 Grenoble, France

[‡] ICBMS, University of Lyon, 43 Bvd. du 11 novembre 1918, 69622, Villeurbanne Cedex, France

ABSTRACT: Bacterial and fungal pathogens involved in lung infection in cystic fibrosis patients utilize a particular family of glycan-binding proteins, characterized by the presentation of six fucose-binding sites on a ring shape scaffold. These lectins are attractive targets for anti-infectious compounds that could interfere in the recognition of host tissues by pathogens. The design of a cyclopeptide-based hexavalent structure allowed for the presentation of six fucose residues. The synthetic hexavalent compound displays appropriate geometry resulting in high avidity binding by lectins from *Aspergillus fumigatus* and *Burkholderia ambifaria*. Replacing the fucose residue with a conformationally constrained fucomimetic does not alter the affinity and provide fine specificity, with no binding to other fucose-specific lectins.

INTRODUCTION

Many microorganisms use lectins as adhesins to interact with host glycoconjugates and to trigger the first step of infection. Among carbohydrate epitopes, fucose has a prominent role since it is an ubiquitous ligand that can be the target of microorganisms receptors on human tissue,² but also in plant cell walls.³ Most opportunistic pathogens are indeed able to spread from soil, to plants and humans. In mammals' tissues, fucose is a marker of inflammation⁴ and fucosylated glycoconjugates are overexpressed in airways of cystic fibrosis patients.⁵

Pathogens responsible for lung infection in those patients, produce soluble lectins which are involved in adhesion of pathogens to fucosylated glycoconjugates such as human histo-blood group epitopes.⁶ Soluble fucose-binding lectins have been characterized in *Pseudomonas aeruginosa*,⁷ *Burkholderia ambifaria*⁸ and airborne fungus *Aspergillus fumigatus*.⁹ Among them, one structural family is of high interest: *B. ambifaria* BambL and *A. fumigatus* AFL (FleA) are made by the repeat of similar small β -sheets of approx. 40 amino acids (Figure 1, panels A and B). Both lectins are able to bind 6 fucose residues with high affinity.^{8,9} The AFL sequence is longer, consisting in six repeats in tandem⁹ while BambL is formed by trimerisation of a two-repeat peptides.¹⁰ In this family, the affinity for fucose is in the micromolar range, an unusually strong binding for protein–carbohydrate interaction.




Figure 1. Crystal structures of (A) BambL and (B) AFL, complexed with fucose or fucoside (PDB codes 4AHA and 3ZW2). (C) Model of compound **1** in intramonomeric site of BambL with amino acids of interest. Hydrogen bonds are represented as blue dotted lines.

In order to hamper pathogen adhesion to host cells, fucose mimetics become attracting tools to design synthetic ligands to intercept fucose-binding bacteria. Monomeric fucoside derivatives have been tested against AFL demonstrating that (2E)-hexenyl α -L-fucopyranoside is a potent functional inhibitor of FleA.¹⁰ With respect to native glycans, glycomimetics are designed to be more resistant to glycosidases, to present a higher bioavailability and to be tailored to decorate scaffolds of different chemical constitutions.^{11, 12}

In this context, we recently reported the synthesis of a fucose-based glycomimetic **1** (Figure 2) presenting a non-natural aryl α -O-fucosyl linkage.¹³ The fucosyl derivative **1** binds to BamBL lectin with an affinity comparable to that of the natural ligand since its fucoside moiety establishes the same contact with BamBL active site as the natural ligand (Figure 1C). Moreover, it presents an unprecedentedly described specificity for BamBL binding site and does not bind to other fucose-specific lectins such as *P. aeruginosa* LecB or human DC-SIGN.




Figure 2. Structure of the aryl fucose mimetic **1**, the hexavalent conjugate **2**, the aminoxyylated fucoside **3** and of the hexavalent conjugate **4**.

Because lectins are multimeric, higher affinity can be obtained by synthesizing multivalent glycoconjugates.^{14, 15} For example, calixarene-based fucosylated clusters display strong affinity for LecB (45 nM) and provide an almost complete protection against *P. aeruginosa* in a mice infection model.¹⁶ Less efforts have been devoted to competing with β -propeller lectins. Fucosylated glycocluster exhibiting 4, 6 or 8 residues were synthesized based on mannose-centered and branched-phosphodiester scaffolds with best affinities of approximately 43 nM.¹⁷ Additionally, hexavalent cyclotrimeratrylene¹⁸ and penta- and decavalent pillarenes^{19, 20} have also

shown very strong affinity for BamBL (ca. 20 nM). Multivalent compounds obtained by dynamic assembly in the presence of BamBL also blocked the entry in epithelial cells with high efficiency.²¹


The selectivity and the simple synthesis of ligand **1** motivated our interest in synthesizing a multivalent system containing the fucose mimetic **1**. This compound was diastereoselectively obtained relying on an efficient inverse electron-demand Diels-Alder reaction. It is stable in the 4-8 pH range and presents a carboxylic hook to introduce a spacer and/or to be linked to a scaffold. To achieve this purpose, we have selected cyclopeptide-based scaffolds which have been proved useful for the construction of tetra-,²² hexa-²³ and hexadecavalent^{24, 25} glycoconjugates with nanomolar affinity for diverse vegetal and bacterial lectins. Herein we report the synthesis of a cyclopeptide-based hexavalent structure displaying fucose mimetic **1** (conjugate **2**) and the binding studies with BamBL, AFL and LecB by isothermal titration calorimetry (ITC). For comparison, the analogue **4**, conjugated with aminooxylated fucose **3** has also been prepared, as previously described.^{26, 27}

RESULTS AND DISCUSSION

Because the target lectins contain six binding sites, we have synthesized two cyclopeptide scaffolds containing 14 amino acid residues (Scheme 1) instead of the usual cyclodecapeptide as previously described.^{28, 29} In a recent study, we have indeed demonstrated that the expansion of the cycle does not disturb the conformation stability of the scaffold which still shows a characteristic β -sheet profile as proved by circular dichroism.²³ The cyclopeptide **5** has been synthesized on solid-phase following a previously described procedure.²³ After Boc-deprotection of **5** upon treatment with TFA, the Boc-NH-PEG-CO₂H linker has been conjugated by standard amide coupling with PyBOP to the lysine side chains and the Boc groups have been successively removed by treatment with TFA to provide scaffold **6**. Hexavalent compound **2** was next prepared from **6** by PyBOP-promoted coupling with fucose mimetic **1**. The reaction occurred quantitatively

at room temperature in 30 min and provided **2** in 50% yield after semi-preparative HPLC purification. Finally, conjugate **4** was synthesized following an oxime strategy. Boc-protected serine residues were first conjugated to **6** then successively deprotected and treated with sodium periodate to generate oxo-aldehyde functions. The resulting compound was finally ligated with aminoxy fucose **3**^{23, 24} in water with 0.1% TFA (pH 2.2) to provide the oxime-linked compound **4** in quantitative yield after purification by HPLC.

Monovalent compound **1** was previously demonstrated to be efficiently bound by BambL with affinity in the same order than the one measured for α -methyl-L-fucoside (α MeFuc) with a K_d close to 1 μ M.¹³ In the present work, we tested the efficiency of compound **1** to bind AFL and compared it to α MeFuc.



Scheme 1. Reagents and conditions: a) i: TFA/CH₂Cl₂/H₂O (47.5/47.5/5), rt, 2h; ii: Boc-NH-PEG-CO₂H, PyBOP, DIPEA, DMF, rt, 30 min; iii: TFA/CH₂Cl₂/H₂O (47.5/47.5/5), rt, 2h, **6** (69% over three steps); b) 1, PyBOP, DIPEA, DMF, rt, 30 min, **2** (50%); c) i: BocSer(tBu)OH, PyBOP, DIPEA, DMF, rt, 2 h; ii: TFA/TIS/H₂O (95/2.5/2.5), rt, 2h; iii: NaIO₄, H₂O, rt, 30 min; iv: **3**, H₂O, 0.1% TFA, **4** (> 90%).

Table 1. Microcalorimetry data for monovalent compounds interacting with BambL and AFL

Compound	Lectin	K_d (μ M)	ΔG (kJ/mol)	ΔH (kJ/mol)	$-T\Delta S$ (kJ/mol)
1	BambL ^a	1.54	-33.2	-28.9	-4.3
	AFL	10.3	-28.5	-28.7	0.2
α MeFuc	BambL ^a	0.99	-34.3	-50.1	15.8
	AFL	43.3	-24.9	-39.2	14.3

^a from⁸

Compound **1** was previously demonstrated to not bind to bacterial lectin LecB and human lectin DC-SIGN.¹³ The precise specificity of **1** for Bambl and AFL makes it a specific inhibitor for the fucose-binding site present in this family of β -propeller lectin. When analyzing the thermodynamics contribution, the binding mode of **1** does not present the entropy barrier, classically observed in protein-carbohydrate interactions. AFL and Bambl interaction with α MeFuc is enthalpy driven with an unfavorable entropy term, while binding to compound **1** does not involve any significant entropy. Multivalent compounds **2** and **4** were evaluated in titration microcalorimetry experiments to determine the stoichiometry and thermodynamic parameters of their binding with LecB, Bambl and AFL (Figure 3, Table 2). Compounds **1** and α MeFuc were used as monovalent references to calculate the relative potency of **2** and **4** respectively. Titrations were performed in the direct injection mode (*i.e.* ligand in syringe and protein in cell).




Figure 3. Isothermal titration calorimetry data: thermograms and corresponding titration curves obtained by injections of compounds **2** and **4** into solutions of BambL (left) and AFL (right). Molar ratio is defined as the number of glycocluster molecules per lectin monomer.

The analysis of the thermodynamic contributions showed that for both compounds, binding is driven by enthalpy but is counterbalanced by a strong entropy barrier resulting from the loss of flexibility induced upon lectin binding. Interestingly, the selectivity of monovalent compound **1** toward BambL and AFL was still observed with hexavalent compound **2**, indicating that the fucose mimetic cannot bind to LecB even at high local concentration. On the contrary, hexafucosylated compound **4** exhibits a K_d of 165 nM for LecB, which confirms that the lack of binding of **2** does not come from an unfavorable geometry of the scaffold.

Table 2. Microcalorimetry data for multivalent compounds interacting with LecB, BambL and AFL

Compound	Lectin	K_d (nM)	N^a	ΔG (kJ/mol)	ΔH (kJ/mol)	$T^*\Delta S$ (kJ/mol)	α
	LecB			No binding			
2	BambL	13.8±3.8	1.01±0.06	-45	-162.5±12.5	-117.5	112
	AFL	44±10	0.68±0.01	-42.1	-115.5±0.5	-73.7	234
4	LecB	165±5	0.60±0.01	-38.7	-133.5±5.5	-94.6	2.6
	BambL	16.8±2.2	1.53±0.1	-44.5	-165±3	-120.5	59.1
	AFL	18.5±8.5	0.77±0.01	-44.4	-154.5±3.5	-109.5	1886 ^l

^a The stoichiometry has been calculated as ratio of multivalent compound for the whole lectin (four binding sites for LecB, six fucose binding sites for trimer of BambL and monomer of AFL). Coefficient α is the relative potency of the glycocluster compared to its monomeric epitope

ITC measurements also showed that hexavalent compounds **2** and **4** bind very strongly to both BambL and AFL. Surprisingly, they both exhibit similar K_d values despite their different epitopes. Compound **4** shows a K_d of 18.5 nM on AFL making it 1886 times more potent than

α MeFuc meaning that each sugar is 314 times more efficient than the monovalent reference. Studies with BambL show that compound **2** binds with a K_d of 13.8 nM resulting in an improvement factor of 112 when compared to **1** and a relative potency of 19 per sugar residue. Stoichiometries for all measurements show an approximate ratio of 1:1 and an enthalpy of binding that is approx. four times stronger than the one measured for α -methyl-fucoside (-162 kJ/mol versus -47 (see Ref 8)) This would indicate that at least four of the six fucose residues are involved in lectin binding. This strong enthalpy is counterbalanced by an important entropy factor necessary to accommodate the clusters' flexible arms in the appropriate direction. Altogether these data indicate that compounds **2** and **4** are the best known ligands for these lectins BambL and AFL to date.




Figure 4. **A.** Model of compound **4** built with extended conformation of linkers. The cyclopeptide was built by extended a related model (see Ref. 30) obtained from NMR studies and molecular modeling. **B.** Model of BambL with compound **1** in the six fucose binding sites (adapted from Ref. 13). **C.** Superimposition of the two models with same scale for size evaluation.

In view of the large increase of affinity observed for hexavalent ligands, molecular modeling was performed in order to evaluate if a single hexameric compound could reach the six binding site of one BambL trimer. As displayed in Figure 4, glycocluster **4** in extended conformation

brings the fucose-analog extremities at distances that are compatible with the fucose binding site on BambL. However, the orientation of the fucose mimetics does not bring their extremities in a favorable location compared to the geometry of the glycoclusters. It is therefore unlikely that the six fucose-analogs would bind at the same time to the same BambL and cross-linking to different proteins is probably occurring.

We describe for the first time glycomimetics with nM affinity for lectins from pathogen organisms, and with precise specificity towards a class of lectins. At the present, the six-bladed fucose-specific propeller that is targeted by these compounds has been described in two human pathogenic microorganisms, i.e. *B. ambifaria* and *A. fumigatus*. The lectin has also been characterized from organisms that interact with plants cell wall, such as RSL from bacterium *Ralstonia solanacearum*, AOL from fungus *Aspergillus oryzae*, both plant pathogens, and AAL mushroom *Aleuria aurantia*. We previously demonstrated that RSL binds to α Fuc1,2Gal disaccharide, an epitope present in both human blood group oligosaccharides, and plant cell wall xyloglucan. Such soluble lectins could be therefore involved in adhesion to different substrates.

B. ambifaria and *A. fumigatus* are both lung pathogens that cause serious infections in immunosuppressed patients, and those with cystic fibrosis. In order to identify other possible target for glycomimetic compounds mimicking fucose, a search based on BambL sequence has been performed in NIH Microbial database. Sequences with similar length and strong identity scores could be identified in several bacteria, some of them being considered as emergent pathogens (Figure 4).

```

B. amb_1 ---MQTAAISWGTT--PSIRVYTAN-GNKITE1RCY2DGS-NWYTGA3FNQ
B. stab_1 ---MQTAAISWGTT--PSIRVYTAN-GNKITE1RCY2DGA-NWYTGA3FNQ
B. pyr_1 ---MQTAAISWGTT--PSIRVYTAN-GNKITE1RCY2DGS-NWYTGA3FAQ
B. cep_1 ---MQTAAISWGTT--PSIRVYTAN-GNKITE1RCY2DGQ-NWYTGA3FNQ
R. pic_1 MSNVQTAATSWGTV--PSIRVYTAN-GGKITE1RCY2DGK-GWYTGA3FNE
R. man_1 MSNVQTAATSWGTV--PSIRVYTAN-GGKITE1RCY2DGK-GWYTGA3FNE
A. fum_1 LFRGTGIAAVNSTN----HLRVYFQDVYGSIR1ESLY2EG--SWANGTEKNVIGNA

B. amb_2 AGDNVSATCWL1SGSAVHIRVYAT-SGGSTTE1WCWDGD-GWTRGAYTGL
B. stab_2 AGDNVSATCWL1SGSAVHIRVYAT-SGGSTTE1YCWDGD-GWTRGGYTGS
B. pyr_2 AGDNVSATCWL1SGSAVHIRVYAT-SGGSTTE1YCWDGD-GWTRGGYTGS
B. cep_2 AGDNVSATCWL1SGSAIHIRVYAT-SGGSTTE1YCWDGD-GWTRGGYTGS
R. pic_2 PGDNVSVTSWL1SGSAIHIRVYAT-SGSNTTE1WCWDGN-SWTKGGYTATN
R. man_2 PGDSVSVTSWL1SGSAIHIRVYAT-SGGSTTE1WCWDGN-SWTKGGYTATN
A. fum_2 KLGSPVAATSKE1L---KHIRVYTLTEGNTLQEFAY2DSGTGWYNGGLGGAK

```

Figure 4. Multiple sequence alignment of BambL related sequences identified in the genomes of Burkholderia and Ralstonia pathogenic species. The two blocks correspond to blade repeats and the four amino acids directly involved in fucose binding are highlighted. Accession number are BAX62607.1 for *B. stabilis*, WP_034182107.1 for *B. pyrrocinia*, WP_059237490.1 for *B. cepacia*, WP_024976975.1 for *R. picketti* and WP_063392503.1 for *R. mannitolilytica*. Sequences of the two first blades of AFL (*A.fum*, amino acids 10 to 103) are included for comparison.

B. stabilis, *B. pyrrocinia*, and *B. cepacia* are members of BCC, the *Burkholderia cepacia* complex, a group of closely related species that are found in various environments, including hospitals, responsible for grave infection in cystic fibrosis patients, and often multi-resistant to antibiotics.³¹ *R. picketti* and *R. mannitolilytica* are also emergent pathogens, now causing more frequent and more severe respiratory infections in CF patients, with most *Ralstonia* identified strains being multi-drug resistant.³²

As displayed in Figure 4, the amino acids involved in fucose binding in this family of lectins are conserved in all described pathogens. It can be therefore expected that the glycomimetics described here, and the ones that could be developed from them in the future will be active on several emergent lung pathogens. Strategies for the use of such anti-infectious compounds are getting more urgent since many strains present increasing multi-resistance to antibiotics.

In summary, we report here a convergent approach that brings together the concepts of glycomimetic and of multivalency for reaching both high affinity and high specificity in targeting the interaction between pathogens and human tissues. Such approach was used previously with

well characterized glycomimetics for blocking the interaction of human DC-SIGN with HIV-glycoprotein³³ and the interaction of cholera-toxin with glycolipids.³⁴ Nevertheless, this is the first report for specific targeting of the fucose-specific β -propeller lectin present in both fungal and bacterial pathogens involved in lung infection in immunocompromised patients and cystic fibrosis ones. The affinities for BamBL and AFL of the glycomimetic hexavalent clusters are in the same range as the fucoclusters as evaluated via ITC studies with dissociation constants in the nanomolar range. The increase in affinity in comparison with the monovalent epitope is very high, with a gain of almost 2000 for binding to AFL, indicating that the cyclopeptide raft scaffold provides efficient multivalency for binding to this lectin family. The non-natural aryl α -O-fucosyl linkage of the glycomimetic adopts a constrained flat conformation that could fit only in the crevasse-shaped binding site of this lectin family, resulting in very precise specificity. Since genes coding for highly similar lectin have been identified in the genome of several emergent pathogens, the present study can be the basis of anti-infectious therapy.

EXPERIMENTAL SECTION

General method: Reaction progress was monitored by reverse-phase HPLC-MS using C₁₈ columns, at 1.3 mL min⁻¹ (Nucleosil 120 Å, 3 μ m C₁₈ particles, 30 x 4.6 mm²) with UV monitoring at 214 nm and 250 nm and with ESI-MS in positive mode, using a linear A-B gradient (5 to 40% B in 15 min, buffer A: 0.09% CF₃COOH in water; buffer B: 0.09% CF₃COOH in acetonitrile). Analytical HPLC analyses were performed by using C₁₈ columns and a linear gradient (5 to 40% B in 25 min), with UV monitoring at 214 nm and 250 nm.

Synthesis of compound 2. To a solution of **6** (30 mg, 0.01 mmol) in dry DMF (3 ml), **1**⁹ (22 mg, 0.072 mmol), PyBOP (37 mg, 0.072 mmol) and DIPEA (31 μ l, 0.18 mmol) were added. The mixture was stirred at rt and after 30 min, analytical HPLC found the quantitative conversion of **6** in **2**. The solvent was evaporated and the peptide was precipitated in Et₂O, centrifuged and washed three times with Et₂O. The crude was purified by semi-preparative HPLC (gradient: 5% to 70% B

in 30 min) to obtain **2** (20 mg, 0.005 mmol, 50%). Analytical HPLC: $R_t = 14.5$ min; MALDI-TOF-MS m/z calcd. for $C_{182}H_{264}N_{26}O_{62}S_6^{2+}$ $[M+2H]^{2+}$ 1999.33 found 2000.00.

Synthesis of compound 4. To a solution of **6** (10 mg, 0.003 mmol) in H_2O (2 ml), $NaIO_4$ (38 mg, 0.18 mmol) was added and the mixture was stirred at rt. After 30 min, the mixture was purified by semi-preparative HPLC (gradient: 5% to 40% B in 30 min) to give a crude, which was used in the next reaction without further purification. A solution of crude and **3** (5 mg, 0.027 mmol) was stirred at rt in a mixture of H_2O + 0.1% TFA (2 ml). After 30 min, analytical HPLC found the quantitative conversion of the intermediate in **4**. The mixture was purified by semi-preparative HPLC (gradient: 5% to 100% B in 30 min) to give pure **4** (11 mg, 0.003 mmol, >90%). MALDI-TOF-MS m/z calcd. for $C_{146}H_{246}N_{32}O_{68}^{2+}$ $[M+2H]^{2+}$ 1768.34 found 1768.40.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI:

Isothermal Titration Calorimetry; Synthesis, RP-HPLC and ESI-MS of compounds **2**, **4**, **5'**, **6**, **8** (PDF).

AUTHOR INFORMATION

Corresponding Authors

*E-mail: cristina.nativi@unifi.it

*E-mail: anne.imberty@cermav.cnrs.fr

*E-mail: olivier.renaudet@ujf-grenoble.fr

Notes

The authors declare no competing financial interest.

Keywords: glycoconjugates • biological activity • lectines • antipathogen agents • glycopeptides

ACKNOWLEDGMENTS

This study was supported by CNRS and Université Grenoble Alpes. O.R. acknowledges the European Research Council Consolidator Grant “LEGO” (647938). This work has been supported by the French ANR project Glyco@Alps (ANR-15-IDEX-02) and Labex ARCANE (ANR-11-LABX-003) and association Vaincre la Mucoviscidose. Fondazione Cassa di Risparmio di Firenze is acknowledged for financial support. Authors thank Mr. Alvaro Brittolli (University of Florence) for having provided a sample of compound **1**.

REFERENCES

- (1) Imberty, A., and Varrot, A. (2008) Microbial recognition of human cell surface glycoconjugates. *Curr. Opin. Struct. Biol.* *18*, 567-576.
- (2) Becker, D. J., and Lowe, J. B. (2003) Fucose: biosynthesis and biological function in mammals. *Glycobiology* *13*, 41R-53R.
- (3) Anderson, C. T., Wallace, I. S., and Somerville, C. R. (2012) Metabolic click-labeling with a fucose analog reveals pectin delivery, architecture, and dynamics in Arabidopsis cell walls. *Proc. Natl. Acad. Sci. U. S. A.* *109*, 1329-34.
- (4) Gringhuis, S., Kaptein, T.M., Wevers, B.A., Mesman, A.W. and Geijtenbeek, T.B. (2014) Fucose-specific DC-SIGN signalling directs T helper cell type-2 responses via IKK ϵ and CYLD-dependent Bcl3 activation. *Nat. Commun.* *5*:3898.
- (5) Glick, M. C., Kothari, V. A., Liu, A., Stoykova, L. I., and Scanlin, T. F. (2001) Activity of fucosyltransferases and altered glycosylation in cystic fibrosis airway epithelial cells. *Biochimie* *83*, 743-7.
- (6) Heggelund, J. E., Varrot, A., Imberty, A., and Krenzel, U. (2017) Histo-blood group antigens as mediators of infections. *Curr. Opin. Struct. Biol.* *44*, 190–200.
- (7) Garber, N., Guempel, U., Gilboa-Garber, N., and Doyle, R. J. (1987) Specificity of the fucose-binding lectin of *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.* *48*, 331-334.
- (8) Audfray, A., Claudinon, J., Abounit, S., Ruvoën-Clouet, N., Larson, G., Smith, D. F., Wimmerová, M., Le Pendu, J., Römer, W., Varrot, A., and Imberty, A. (2012) The fucose-binding lectin from opportunistic pathogen *Burkholderia ambifaria* binds to both plant and human oligosaccharidic epitopes. *J. Biol. Chem.* *287*, 4335-4347.
- (9) Houser, J., Komarek, J., Kostlanova, N., Cioci, G., Varrot, A., Kerr, S. C., Lahmann, M., Balloy, V., Fahy, J. V., Chignard, M., Imberty, A., and Wimmerova, M. (2013) A soluble fucose-specific lectin from *Aspergillus fumigatus* conidia - Structure, specificity and possible role in fungal pathogenicity. *PLoS ONE* *8*, e83077.
- (10) Kerr, S. C., Fischer, G. J., Sinha, M., McCabe, O., Palmer, J. M., Choera, T., Lim, F. Y., Wimmerova, M., Carrington, S. D., Yuan, S., Lowell, C. A., Oscarson, S., Keller, N. P., and Fahy, J. V. (2016) FleA Expression in *Aspergillus fumigatus* Is Recognized by

- Fucosylated Structures on Mucins and Macrophages to Prevent Lung Infection. *PLoS Pathogens* 12, e1005555.
- (11) Cecioni, S., Imberty, A., and Vidal, S. (2015) Glycomimetics versus multivalent glycoconjugates for the design of high affinity lectin ligands. *Chem. Rev.* 115, 525-61.
 - (12) Ernst, B., and Magnani, J. L. (2009) From carbohydrate leads to glycomimetic drugs. *Nat. Rev. Drug Discov.* 8, 661-77.
 - (13) Richichi, B., Imberty, A., Gillon, E., Bosco, R., Sutkeviciute, I., Fieschi, F., and Nativi, C. (2013) Synthesis of a selective inhibitor of a fucose binding bacterial lectin from *Burkholderia ambifaria*. *Org. Biomol. Chem.* 11, 4086-4094.
 - (14) Chabre, Y. M., Giguère, D., Blanchard, B., Rodrigue, J., Rocheleau, S., Neault, M., Rauthu, S., Papadopoulos, A., Arnold, A., Imberty, A., and Roy, R. (2011) Combining glycomimetic and multivalent strategies toward designing potent bacterial lectin inhibitors. *Chem. Eur. J.* 17, 6545-6562.
 - (15) Bernardi, A., Jiménez-Barbero, J., Casnati, A., Castro, C. D., Darbre, T., Fieschi, F., Finne, J., Funken, H., Jaeger, K.-E., Lahmann, M., Lindhorst, T. K., Marradi, M., Messner, P., Molinaro, A., Murphy, P., Nativi, C., Oscarson, S., Penadés, S., Peri, F., Pieters, R. J., Renaudet, O., Reymond, J.-L., Richichi, B., Rojo, J., Sansone, F., Schäffer, C., Turnbull, W. B., Velasco-Torrijos, T., Vidal, S., Vincent, S., Wennekes, T., Zuilhof, H., and Imberty, A. (2013) Multivalent glycoconjugates as anti-pathogenic agents. *Chem. Soc. Rev.* 42, 4709 - 4727.
 - (16) Boukerb, A., Rousset, A., Galanos, N., Méar, J.-B., Thepaut, M., Grandjean, T., Gillon, E., Cecioni, S., Abderrahmen, C., Faure, K., Redelberger, D., Kipnis, E., Dessein, R., Havet, S., Darblade, B., Matthews, S., de Bentzmann, S., Guéry, B., Cournoyer, B., Imberty, A., and Vidal, S. (2014) Anti-adhesive properties of glycoclusters against *Pseudomonas aeruginosa* lung infection. *J. Med. Chem.* 57, 10275-10289.
 - (17) Ligeour, C., Audfray, A., Gillon, E., Meyer, A., Galanos, N., S.Vidal, Vasseur, J. J., Imberty, A., and Morvan, F. (2013) Synthesis of antenna and mannose-centered fucosylated glycoclusters and their binding studies with *Burkholderia ambifaria* lectin (BambL). *RSC Adv.* 3, 19515-19524.
 - (18) Galanos, N., Chen, Y., Zachary, P. M., Gillon, E., Dutasta, J. P., Star, A., Imberty, A., Martinez, A., and Vidal, S. (2016) Cyclotrimeratrylene-based glycoclusters as high affinity ligands of bacterial lectins from *Pseudomonas aeruginosa* and *Burkholderia ambifaria*. *ChemistrySelect* 1, 5863-5868
 - (19) Buffet, K., Nierengarten, I., Galanos, N., Gillon, E., Holler, M., Imberty, A., Matthews, S. E., Vidal, S., Vincent, S. P., and Nierengarten, J. F. (2016) Pillar[5]arene-based glycoclusters: synthesis and multivalent binding to pathogenic bacterial lectins. *Chem. Eur. J.* 22, 2955 – 2963.
 - (20) Galanos, N., Gillon, E., Imberty, A., Matthews, S. E., and Vidal, S. (2016) Pentavalent pillar[5]arene-based glycoclusters and their multivalent binding to pathogenic bacterial lectins. *Org. Biomol. Chem.* 14, 3476-3481.
 - (21) Machida, T., Novoa, A., Gillon, E., Zheng, S., Claudinon, J., Eierhoff, T., Imberty, A., Romer, W., and Winssinger, N. (2017) Dynamic cooperative glycan assembly blocks the binding of bacterial lectins to epithelial cells. *Angew. Chem. Int. Ed.* 56, 6762-6766.

- (22) Fiore, M., Berthet, N., Marra, A., Gillon, E., Dumy, P., Dondoni, A., Imberty, A., and Renaudet, O. (2013) Tetravalent glycocyclopeptide with nanomolar affinity to wheat germ agglutinin *Org. Biomol. Chem.* *11*, 7113-7122.
- (23) Pifferi, C., Goyard, D., Gillon, E., Imberty, A., and Renaudet, O. (2017) Synthesis of mannosylated glycodendrimers and evaluation against BC2L-A lectin from *Burkholderia cenocepacia*. *ChemPlusChem* *82*, 390 – 398
- (24) Berthet, N., Thomas, B., Bossu, I., Dufour, E., Gillon, E., Garcia, J., Spinelli, N., Imberty, A., Dumy, P., and Renaudet, O. (2013) High affinity glycodendrimers for the lectin LecB from *Pseudomonas aeruginosa*. *Bioconjug. Chem.* *24*, 1598-1611.
- (25) Thomas, B., Pifferi, C., Daskhan, G. C., Fiore, M., Berthet, N., and Renaudet, O. (2015) Divergent and convergent synthesis of GalNAc-conjugated dendrimers using dual orthogonal ligations. *Org. Biomol. Chem.* *13*, 11529-38.
- (26) Bossu, I., Richichi, B., Dumy, P., and Renaudet, O. (2011) Synthesis of O-(6-deoxy- α - and β -L-galactopyranosyl) hydroxylamines (α - and β -L-fucopyranosylhydroxylamines), in *Carbohydrate Chemistry: Proven Synthetic Methods* (Kovac, P., Ed.) pp 377–386, CRC Press Boca Raton, FL.
- (27) Dulery, V., Renaudet, O., Philouze, C., and Dumy, P. (2007) Alpha and beta L-Fucopyranosyl oxyamines: key intermediates for the preparation of fucose-containing glycoconjugates by oxime ligation. *Carbohydr. Res.* *342*, 894-900.
- (28) Daskhan, G. C., Berthet, N., Thomas, B., Fiore, M., and Renaudet, O. (2015) Multivalent glycocyclopeptides: toward nano-sized glycostructures. *Carbohydr. Res.* *405*, 13-22.
- (29) Galan, M. C., Dumy, P., and Renaudet, O. (2013) Multivalent glyco(cyclo)peptides. *Chem. Soc. Rev.* *42*, 4599-612.
- (30) Bossu, I., Šulc, M., Křenek, K., Dufour, E., Garcia, J., Berthet, N., Dumy, P., Křen, V., and Renaudet, O. (2011) Dendri-RAFTs: a second generation of cyclopeptide-based glycoclusters. *Org. Biomol. Chem.* *9*, 1948-59.
- (31) Vandamme, P., and Dawyndt, P. (2011) Classification and identification of the *Burkholderia cepacia* complex: Past, present and future. *Syst. Appl. Microbiol.* *34*, 87-95.
- (32) Lipuma, J. J. (2010) The changing microbial epidemiology in cystic fibrosis. *Clin. Microbiol. Rev.* *23*, 299-323.
- (33) Sattin, S., Daghetti, A., Thepaut, M., Berzi, A., Sanchez-Navarro, M., Tabarani, G., Rojo, J., Fieschi, F., Clerici, M., and Bernardi, A. (2010) Inhibition of DC-SIGN-mediated HIV infection by a linear trimannoside mimic in a tetravalent presentation. *ACS Chem. Biol.* *5*, 301-12.
- (34) Arosio, D., Fontanella, M., Baldini, L., Mauri, L., Bernardi, A., Casnati, A., Sansone, F., and Ungaro, R. (2005) A synthetic divalent cholera toxin glycoconjugate having higher affinity than natural GM1 oligosaccharide. *J. Am. Chem. Soc.* *127*, 3660-1.