



HAL
open science

Phospholipase D inhibitors screening: Probing and evaluation of ancient and novel molecules

Yani Arhab, Karim Bessaa, Houda Abla, Meryem Aydin, Renaud Rahier, Arnaud Comte, Leyre Brizuela, Saïda Mebarek, Florent Perret, Mickaël Cherrier, et al.

► **To cite this version:**

Yani Arhab, Karim Bessaa, Houda Abla, Meryem Aydin, Renaud Rahier, et al.. Phospholipase D inhibitors screening: Probing and evaluation of ancient and novel molecules. *International Journal of Biological Macromolecules*, 2021, 166, pp.1131-1140. 10.1016/j.ijbiomac.2020.10.268 . hal-03039714

HAL Id: hal-03039714

<https://hal.univ-grenoble-alpes.fr/hal-03039714v1>

Submitted on 27 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

International Journal of Biological Macromolecules

Phospholipase D inhibitors screening: probing and evaluation of ancient and novel molecules

--Manuscript Draft--

Manuscript Number:	IJBIMAC-D-20-01603R2
Article Type:	Research Paper
Section/Category:	Proteins and Nucleic acids
Keywords:	phospholipase D, inhibitor, phosphatidic acid
Corresponding Author:	Alexandre Noiriél Univ Lyon, Université Lyon 1, Institut de Chimie et de Biochimie Moléculaires et Supramoléculaires (ICBMS), UMR 5246 CNRS, Métabolisme, Enzymes et Mécanismes Moléculaires (MEM2) Villeurbanne, FRANCE
First Author:	Yani Arhab
Order of Authors:	Yani Arhab Karim Bessaa Houda Abla Meryem Aydin Renaud Rahier Arnaud Comte Leyre Brizuela Saida Mebarek Florent Perret Mickael Cherrier Abdelkarim Abousalham Alexandre Noiriél
Abstract:	Phospholipase D (PLD) is a ubiquitous enzyme that cleaves the distal phosphoester bond of phospholipids generating phosphatidic acid (PA). In plants, PA is involved in numerous cell responses triggered by stress. Similarly, in mammals, PA is also a second messenger involved in tumorigenesis. PLD is nowadays considered as a therapeutic target and blocking its activity with specific inhibitors constitutes a promising strategy to treat cancers. Starting from already described PLD inhibitors, this study aims to investigate the effect of their structural modifications on the enzyme's activity, as well as identifying new potent inhibitors of eukaryotic PLDs. Being able to purify the plant PLD from <i>Vigna unguiculata</i> (VuPLD), we obtained a SAXS model of its structure. We then used a fluorescence-based test suitable for high-throughput screening to review the effect of eukaryotic PLD inhibitors described in the literature. In this regard, we found that only few molecules were in fact able to inhibit VuPLD and we confirmed that vanadate is the most potent of all with an IC50 around 58 µM. Moreover, the small-scale screening of a chemical library of 3,120 compounds allowed us to optimize the different screening's steps and paved the way towards the discovery of new potent inhibitors.
Suggested Reviewers:	Michael Frohman Michael.Frohman@Stonybrook.edu Johanna Mansfeld mansfeld@biochemtech.uni-halle.de Martin-Luther University Halle-Wittenberg, Institute of Biochemistry and Biotechnology, Kurt-Mothes-Str. 3, 06120 Halle (Germany).

	<p>Sona Spandey spandey@danforthcenter.org Dr. Sona Pandey, Donald Danforth Plant Science Center, 975 N. Warson Rd., St. Louis, MO 63132 (USA)</p>
	<p>Dorothea Bartels dbartels@uni-bonn.de Pr. Dorothea Bartels, Institute of Molecular Physiology and Biotechnology of Planta (IMBIO), University of Bonn Kirschallee 1, 53115 Bonn (Germany)</p>
Opposed Reviewers:	
Response to Reviewers:	

Phospholipase D inhibitors screening: probing and evaluation of ancient and novel molecules

Yani Arhab^{‡a}, *Karim Bessaa*^{‡a}, *Houda Abla*^{‡a}, *Meryem Aydin*[‡], *Renaud Rahier*[‡], *Arnaud Comte*[†], *Leyre Brizuela*[‡], *Saïda Mebarek*[‡], *Florent Perret*⁺, *Mickaël V. Cherrier*[†], *Abdelkarim Abousalham*[‡], and *Alexandre Noiriel*^{‡*}

[‡] Univ Lyon, Université Lyon 1, Institut de Chimie et de Biochimie Moléculaires et Supramoléculaires (ICBMS), UMR 5246 CNRS, Métabolisme, Enzymes et Mécanismes Moléculaires (MEM²), Bât Raulin, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne cedex, France.

[†] Univ Lyon, Université Lyon 1, Institut de Chimie et de Biochimie Moléculaires et Supramoléculaires (ICBMS), UMR 5246 CNRS, Chimiothèque, Bât Lederer, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne cedex, France.

⁺ Univ Lyon, Université Lyon 1, Institut de Chimie et de Biochimie Moléculaires et Supramoléculaires (ICBMS), UMR 5246 CNRS, Chimie Supramoléculaire Appliquée (CSAp), Bât Raulin, 43 Bd du 11 Novembre 1918, F-69622 Villeurbanne cedex, France.

[†] Univ. Grenoble Alpes, CEA, CNRS, IBS, Metalloproteins, F-38000 Grenoble, France.

^a These authors contributed equally to this work.

* To whom correspondence should be addressed

Keywords: phospholipase D, inhibitor, phosphatidic acid.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

21 **Abstract**

22 Phospholipase D (PLD) is a ubiquitous enzyme that cleaves the distal phosphoester bond of
23 phospholipids generating phosphatidic acid (PA). In plants, PA is involved in numerous cell
24 responses triggered by stress. Similarly, in mammals, PA is also a second messenger involved in
25 tumorigenesis. PLD is nowadays considered as a therapeutic target and blocking its activity with
26 specific inhibitors constitutes a promising strategy to treat cancers. Starting from already described
27 PLD inhibitors, this study aims to investigate the effect of their structural modifications on the
28 enzyme's activity, as well as identifying new potent inhibitors of eukaryotic PLDs. Being able to
29 purify the plant PLD from *Vigna unguiculata* (VuPLD), we obtained a SAXS model of its
30 structure. We then used a fluorescence-based test suitable for high-throughput screening to review
31 the effect of eukaryotic PLD inhibitors described in the literature. In this regard, we found that
32 only few molecules were in fact able to inhibit VuPLD and we confirmed that vanadate is the most
33 potent of all with an IC₅₀ around 58 μM. Moreover, the small-scale screening of a chemical library
34 of 3,120 compounds allowed us to optimize the different screening's steps and paved the way
35 towards the discovery of new potent inhibitors.

36 Introduction

37 Phospholipase D (PLD) (EC 3.1.4.4) is a diverse family of enzymes found in microbes, fungi,
38 viruses, plants and animals [1]. In addition to its hydrolytic activity towards phospholipids
39 releasing phosphatidic acid (PA) and a polar moiety, PLD catalyzes a transphosphatidylation
40 reaction as well. This reaction occurs when a primary alcohol is used instead of water, making
41 possible the synthesis of rarer phospholipids by exchanging the polar headgroup, as
42 phosphatidylserine from the abundant phosphatidylcholine (PC) in the presence of serine.

43 Contrary to several bacterial PLDs, plant and mammalian PLDs share a common feature
44 represented by a duplicated HKD motif that is involved in the catalytic reaction, there are then two
45 classes of PLDs based on their primary structure: the HKD-containing and the non-HKD [2]. This
46 latter family regroups well-used enzymes that belongs to the superfamily of PLC-like
47 phosphodiesterases [3], such as the commercial PLD from *Streptomyces chromofuscus* [4] or the
48 PLD from arachnids [5]. The HKD-containing PLD family belongs to the PLD superfamily and is
49 mostly found in mammals as for example PLD1 and PLD2 in *Homo sapiens*, in plants as PLD α ,
50 in yeast with SPO14 [2,6,7], and in several microbes as the crystallized PLD from *Streptomyces*
51 *sp.* strain PMF[8]. The hallmark of the PLD superfamily is the presence of the so-called HKD
52 motif that is formed by a conserved His, Lys and Asp in a typical HxKx₄D sequence that was
53 recently refined [9]. In plants and mammals, the HKD motif is duplicated in the primary structure,
54 and both motifs are thought to be closed in the tertiary structure to form the catalytic site at the
55 interface of the bi-lobed enzyme [10–13].

56 Furthermore, one common mechanism of action defined as a two-step “ping-pong” reaction
57 mechanism is conserved between plant and mammalian PLDs [2]. The interest in PLD as a drug
58 target is due to the fact that PLDs are essential to intracellular and extracellular signaling. PLD

1
2
3
4 59 have been described to be involved in settings ranging from platelet activation to response to
5
6
7 60 cardiac ischemia, viral infections, neurodegenerative diseases, and in cancers. Therefore, human
8
9 61 PLD are a valid target for a variety of drug therapies [14].

10
11 62 Our work focuses on identifying potential new inhibitor of eukaryotic PLDs, especially the human
12
13
14 63 isoforms, by using a plant PLD as a model. Indeed, VuPLD shows an overall 19 % peptidic
15
16 64 sequence identity with human PLD1, which reaches 42 % and 69 % around the first and second
17
18
19 65 HKD motif, respectively. Thus, based on these elements, we hypothesized that an inhibitor
20
21 66 directed against the catalytic site of plant PLD would somehow be able to inhibit mammalian PLD
22
23
24 67 and *vice versa*. Previous works to screen inhibitors on mammalian PLD were weakened by the
25
26 68 difficulty to obtain sufficient amounts of enzyme for HTS of PLD inhibitors. This difficulty was
27
28
29 69 partially circumvented, either by the use of the atypical prokaryotic PLD enzyme from
30
31 70 *Streptomyces* for preliminary screenings [15], by using *in silico* methods [16], by virtual screening
32
33
34 71 [17], or other undisclosed conditions [18]. Recombinant plant PLD α is however easy to produce
35
36 72 and could be purified in one step [19]. Therefore, considering their similarities at multiple levels,
37
38
39 73 we assumed that a molecule capable of inhibiting plant PLDs could as well affect human PLDs.
40
41 74 This strategy could thus constitute an interesting and original approach to identify novel inhibitors
42
43 75 than using non-HKD PLDs as templates for the screening.

44
45 76 Only a few inhibitors of PLDs were described in the literature. N-acylethanolamines (NAEs) are
46
47
48 77 a class of potent lipids released from the membrane phospholipid N-acylphosphatidylethanolamine
49
50
51 78 (NAPE), and are found as endogenous constituents of desiccated seeds. The hydrolysis of NAPE
52
53 79 is catalyzed by a NAPE-specific PLD or PLD-like enzyme [20], but not by canonical PLD such as
54
55 80 plant PLD α . However, NAE is able to inhibit PLD α activity with an IC₅₀ estimated at 0.15 μ M as
56
57
58 81 for NAE (12:0) [21]. Applying NAE (12:0) improved seed desiccation tolerance via the inhibition
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

82 of PLD α , that led to the reduction of PA formation, PA being a key modulator of desiccation
83 sensitivity [22]. Likewise, lysophosphatidylethanolamine (LPE) was shown to be a specific
84 inhibitor of plant PLDs, that could act as a lipid-derived senescence retardant via the inhibition of
85 PLD [23]. As an example adding 10 μ M of LPE (18:1) reduced by half the activity of PLD partially
86 purified from cabbage [23].

87 Similarly, the immunosuppressant ciclosporin at 5 μ M was also shown to completely inhibit the
88 cabbage PLD activity [24], and so were alkylphosphate esters [25], octadecylphosphocholine
89 being the most efficient inhibitor of this class of molecules with an IC₅₀ of 6.4 μ M. However, these
90 molecules having a classical phosphoester bond they are substrate as well of PLD enzymes [26].
91 In fact, compared to PC, the canonical substrate tested in the same conditions, around 3% of the
92 specific activity was found when hexadecylphosphocholine was used [25]. Hexanal and its reduced
93 form hexanol have been shown to be effective PLD inhibitors in different subcellular fractions
94 from corn kernels, and PLD activity was inhibited from 50 to 70 % using 0.1 % of hexanal in *in*
95 *vitro* assays [27,28]. Volatile hexanal being a natural product of the catabolism of linoleic and
96 linolenic acid during senescence, it was used as a check-point for arresting membrane lipid
97 degradation in plants, and consequently enhances shelf-life of fruits as described in raspberry [29]
98 and in litchi [30]. PLD inhibition is used in banana when treated with hexanal, to thicken the cell
99 wall and impeded penetration of pathogenic spores [31]. Moreover, PLD in rice causes rancidity and
100 'stale flavor' in bran oil, and thus limits the rice bran usage [32], demonstrating the need for specific
101 plant PLD inhibitors.

102 As far as mammalian PLDs are concerned, several compounds were reported in literature. For
103 instance, fodrin [33] indirectly inhibits the PLD activity by decreasing the amount of
104 phosphatidylinositol (4,5)-bisphosphate (PIP₂), a cofactor necessary for mammalian PLD activity.

1
2
3
4 105 However, this compound is not considered useful for research or clinical trials. Halopemide was
5
6 106 identified by HTS in 2007 as a modest inhibitor of PLD2 [18]. Several variants of halopemide
7
8
9 107 were then developed to exhibit a preferential selectivity towards either PLD1 or PLD2 [34]. *In*
10
11 108 *vitro*, halopemide has an IC₅₀ of 220 and 310 nM for PLD1 and PLD2 respectively. Conversely,
12
13
14 109 VU0155069 is more selective towards PLD1, with an IC₅₀ of 46 nM towards PLD1 compared to
15
16 110 933 nM towards PLD2 *in vitro*, and 11 nM and 1800 nM in cells respectively, whereas a
17
18
19 111 VU0364739 precursor is more selective towards PLD2 (IC₅₀ of 140 nM *in vitro* and of 110 nM in
20
21 112 cells, compared to 5.1 μM *in vitro* and 1 μM in cells towards PLD1) [35]. Interestingly 5-fluoro-
22
23
24 113 2-indolyl des-chlorohalopemide (FIPI) which at approximately 25 nM induced a 50% loss of
25
26 114 activity independently on PLD1 and PLD2 [36] is thought to act on the catalytic site of both
27
28
29 115 isoforms PLD1 and PLD2. In the opposite VU0364739 which is selective of PLD2 [37] was
30
31 116 previously shown by computational modelling to use an allosteric site overlapping with the PH
32
33 117 domain to target its activity [38], but surface plasmon resonance experiments have demonstrated
34
35
36 118 that this family of inhibitors binds directly to the catalytic domain [11]. FIPI's IC₅₀ on PLD from
37
38 119 *Streptomyces* strain PMF was reported to be around 100 nM, which is 4 to 6 times higher than the
39
40
41 120 values found on PLD1 and PLD2, demonstrating that some inhibitors directly block enzyme and
42
43 121 phospholipid vesicles substrate binding [15]. Last but not least, resveratrol, an antioxidant found
44
45
46 122 in red wine, was described to inhibit plant PLD [39], and human isoforms as well with an IC₅₀
47
48 123 around 61 μM [40].
49
50 124 In order to find the optimal conditions for the screening and to understand the best features of a
51
52
53 125 PLD inhibitor, we first characterized the purified enzyme VuPLD and described its physical shape
54
55 126 and biochemical structural properties. Then, we characterized the effect of already-known
56
57
58 127 inhibitors of eukaryotic PLD on VuPLD. Several types of inhibitors have been described in the
59
60
61
62
63
64
65

1
2
3
4 128 literature and could be basically divided into three classes. First, phosphate mimicking ions such
5
6
7 129 as vanadate [41] or tungstate that compete with the phosphate moiety of the phospholipid substrate.
8
9 130 Secondly natural products such as resveratrol [39] and ciclosporine [24], that are small organic
10
11
12 131 compounds supposedly protecting enterocyte mitochondria from oxidative stress. Finally,
13
14 132 halopemide and its derivatives that were specially designed for the two isoforms of mammalian
15
16 133 PLD with catalytic activity (PLD1 and PLD2). We then screened a chemical library of roughly
17
18
19 134 3,120 molecules of diverse origins. The optimization of a 6-step screen could now be used for
20
21 135 larger libraries.
22

23
24 136

26 137 **Material and Methods**

28 29 30 138 **Reagents and material**

31
32
33 139 1,2-Dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC), choline oxidase from *Arthrobacter*
34
35 140 *globiformis*, horseradish peroxidase (type VI), bovine serum albumin (BSA), 8-hydroxyquinoline
36
37
38 141 (8HQ), resveratrol, ciclosporin, VU0285655, alcohols, hexanal, N-acylethanolamine, and mineral
39
40 142 salts: sodium vanadate, vanadate derivatives (oxytriethoxide, oxytripropoxide, oxytriisopropoxide,
41
42
43 143 tris(triphenylsiloxy)), tungstate and phosphate came from Sigma Aldrich-Fluka Chimie.
44
45 144 Halopemide, VU0359595, and FIPI were from Cayman Chemical. Palmitoyl-2-oleoyl-*sn*-glycero-3-
46
47
48 145 phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphate (POPA) in chloroform
49
50 146 were purchased from Avanti Polar Lipids (Alabama, USA). Thin layer chromatography (TLC)
51
52
53 147 plates were purchased from Macherey-Nagel GmbH & Co. Kg (Germany). Mammalian-PLDs
54
55 148 inhibitors halopemide and derivatives were from Cayman chemicals. Mineral salts: sodium
56
57
58 149 vanadate and its derivatives, tungstate and phosphate were dissolved in water. All PLD inhibitors
59
60 150 and compounds from the ICBMS chemical library were prepared in DMSO (DMSO levels used
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

151 in this study remained below 5%). Flat-bottom black 96-well microplates (Greiner bio-one) were
152 used for fluorescence measurements.

153
154 Purification of VuPLD
155 Recombinant VuPLD was produced in *Pichia pastoris* and purified as described previously [19].
156 Small aliquots were thawed and diluted before each experiment. A single batch was used for the
157 enzymatic characterization, and another for the screening of the chemical library. Protein
158 concentrations were determined using the Bradford's procedure [42], with Dye Reagent and BSA
159 as the standard.

160
161 Small angle X-ray scattering experiment
162 Data were collected at the ESRF beamline BM29 (Grenoble, France). VuPLD was concentrated
163 at 2.55 mg/mL in PIPES buffer 10 mM, pH 6.2, containing EDTA 0.1 mM and NaCl 200 mM.
164 Reconstruction of the overall shape of the VuPLD from X-ray scattering data was first achieved
165 by the *ab initio* modeling programs DAMMIN (20 models calculated) [43] or GASBOR (50
166 models calculated) [44], then models were averaged using DAMAVER [45]. Comparison of the
167 SAXS data and protein structure (PDB code : 6KZ9) was done using FoXS [46]. Protein molecular
168 mass was estimated from scattering intensities $I(0)$ using SAXSMoW2 software [47].

169
170 PLD activity measurement

1
2
3
4 171 *Chromogenic assay*

5
6
7 172 A detailed protocol of this assay has been recently published [48]. Briefly, 0.4 mM of DMPC in
8
9 173 SDS/Triton-X100 mixed micelles were incubated with pure PLD (0.7 to 0.9 $\mu\text{g}/\text{mL}$, final
10
11 174 concentration) in the presence or absence of inhibitors for 10 min at 30°C. The reaction was started
12
13 175 by the addition of 20 mM of CaCl_2 , then blocked after 10 min by addition of 83.3 mM of EDTA,
14
15 176 and revealed by addition of peroxidase, choline oxidase, 2-hydroxy-3,5-dichlorobenzenesulfonate
16
17 177 and 4-aminoantipyrine. The PLD-generated choline was quantified with end point measurements
18
19 178 by recording the absorbance at 500 nm, based on a standard curve obtained with pure choline.
20
21
22
23
24
25
26
27
28

29 179

30
31 180 *Fluorescence-based assay*

32 181 The preparation of 8HQ / phospholipid/ detergent mixtures was adapted from [49] with minor
33
34 182 modifications to control the activation of the enzymes. To prepare the mixtures, 40 nmol of POPC
35
36 183 (266 μM , final concentration) were first dissolved in chloroform, then evaporated under a stream
37
38 184 of nitrogen. Subsequently, dried phospholipids were dispersed in a buffer containing 50 mM Tris-
39
40 185 HCl, pH 8.0, 1 mM Triton X-100, 0.7 mM SDS, 66 μM 8HQ, and 50 μM EDTA. The micellar
41
42 186 solution was vortexed for 30 s, sonicated for 10 min using a bath sonicator (Deltasonic type O11C,
43
44 187 Fécamp, France), then vortexed for an additional 30 s. One pmol of enzyme (VuPLD) was finally
45
46 188 added as well as the compounds to be tested. After a 10-minute incubation period at 30°C, the
47
48 189 reaction was triggered by adding 20 mM of CaCl_2 . Fluorescence measurements were recorded at
49
50 190 30°C using a microtiter plate fluorescence reader (Tecan Infinite M200, Austria) at wavelengths
51
52 191 366 nm and 490 nm of excitation and emission, respectively. To monitor PLD activity, a baseline
53
54 192 of 4 kinetic cycles of 30 s was first recorded, after which 20 mM CaCl_2 were injected in each well
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

193 to trigger the PLD hydrolysis. Fluorescence intensities were then measured during a 1h-kinetic
194 cycle with interval shaking before each read. The PLD-generated PA was quantified by recording
195 the fluorescence in the same conditions as described above, based on a standard curve obtained
196 with pure POPA [49].

197
198 TLC analysis
199 The phospholipids were extracted with chloroform/methanol (2:1, v/v). The lower organic phase
200 was collected, dried, and subjected to TLC separation together with proper phospholipid standards.
201 TLC plates were developed in a solvent system using a mixture of chloroform/methanol/water
202 (65:25:5, v/v/v) as the eluent. Phospholipid spots were revealed by charring the plate after spraying
203 a mixture of 10 % copper sulfate and 10 % phosphoric acid in water and heating at 150°C for 5
204 min.

206 Statistical analysis
207 Statistically significant differences were determined by Brown-Forsyth and Welsh ANOVA test
208 using the computer program Prism (GraphPad Software). Results were considered significant
209 when $p < 0.05$ (*), highly significant when $p < 0.01$ (**), and extremely significant when $p < 0.001$
210 (***).

212 Results

213 Modelization of VuPLD by SAXS experiment

1
2
3
4 214 VuPLD was produced according to Arhab *et al.*[48]. The amount of homogenous VuPLD obtained
5
6 215 after purification was 1 mg per liter of yeast culture. The specific activity toward DMPC was 20
7
8
9 216 U/mg. Scattering measurement with synchrotron radiation were performed at pH 6.2 with 2.55
10
11 217 mg/mL of VuPLD. A typical SAXS profile of VuPLD is shown in Fig. 1A and a Kratky curve
12
13 218 displays a symmetric peak indicating that VuPLD is globular and correctly folded (Fig. 1B). From
14
15
16 219 scattering intensities $I(0)$ a molecular mass of 89.1 kDa was calculated, which is in accordance
17
18 220 with a monomer of VuPLD as the theoretical molecular mass deduced from the amino acid
19
20
21 221 sequence is 91.5 kDa. The radius of gyration is 2.81 +/- 0.05 nm and D_{\max} (maximum dimension)
22
23 222 is calculated to be 9.89 nm (Fig. 1C). The calculated model by DAMMIN shows a globular and
24
25
26 223 symmetric molecule with extended extremities and an ovoid shape (Fig. 1D). The program
27
28 224 GASBOR calculated a very similar model however this one is less symmetrical and displays a
29
30
31 225 larger base and a fine apical extremity (Fig. S1). The comparison of the collected SAXS data and
32
33 226 calculated scattering curve from the published structure of the PLD show a good correlation (Fig.
34
35
36 227 S2A) indicating that the protein adopts a similar conformation in the crystals and in solution. The
37
38 228 small differences observed could be due to small motion of the C2 domain.
39
40

41 229

42 43 230 **Characterization of the salts effect on PLD activity**

44
45 231 Ions such phosphate, tungstate and vanadate salts were already reported as potent inhibitors of
46
47 232 partially purified plant PLDs. We were able to confirm these results by measuring an IC_{50} for these
48
49
50 233 three ions on a pure VuPLD. Vanadate was the most potent of them with an IC_{50} of 58 μ M and
51
52 234 being mostly able to totally inhibit the enzymatic activity at a concentration of 260 μ M (Fig. 2A)
53
54
55 235 and was consequently used as a reference for HTS of PLD inhibitors. Likewise, tungstate was also
56
57
58 236 able to inhibit totally the PLD activity at a concentration of 2 mM, and harbored an IC_{50} value of
59
60
61
62
63
64
65

1
2
3
4 237 375 μM (Fig. 2C). Phosphate behaved in between, with an IC_{50} estimated to 125 μM but not being
5
6
7 238 able to totally abolish the activity, 25% of residual activity was quantified at 1 mM of phosphate
8
9 239 that is the maximal concentration usable in this test (Fig. 2B), as Ca^{2+} cofactor precipitates at higher
10
11 240 concentration of phosphate.

13
14 241 An unusual slight synergistic and inhibitory effect was observed when phosphate and vanadate
15
16 242 were incubated together (Fig. 2D). Indeed, when both molecules were incubated together at their
17
18
19 243 respective IC_{50} , only 15 % of residual activity was measured that was lower than the theoretical
20
21 244 25 % expected at these particular concentrations, suggesting the combined salts potentiate the PLD
22
23 245 inhibition.

24
25
26 246

27 28 29 247 **Vanadate derived compounds inhibitory effect on PLD activity**

30
31 248 Vanadate being the PLD inhibitor with the more potent inhibitory impacts, we decided to examine
32
33 249 the effects of the environment around this particular ion by testing diverse vanadate derivatives
34
35
36 250 such as vanadium oxytriethoxide, vanadium oxytripropoxide, vanadium oxytriisopropoxide and
37
38 251 tris(triphenylsiloxy) vanadium oxide, as all these molecules harbor different hydrocarbon chain
39
40
41 252 lengths and steric environment (Fig. S3). If vanadate took the place of the phosphate moiety of
42
43 253 phospholipid substrate in the active site, changes of alkyl chain length and steric environment
44
45 254 around vanadate would have been expected to modify its IC_{50} . However, even if the IC_{50} of
46
47
48 255 vanadate derivatives were lower than the one of vanadate (Fig. S4), no dramatic variation could
49
50 256 be observed amongst these four derivatives (Fig. 3). IC_{50} were ranging from 7 μM (vanadate
51
52
53 257 oxytriethoxide) to 15 μM (vanadate oxytriisopropoxide).

54
55 258

56 57 58 259 **Alcohols inhibitory effect on PLD hydrolysis activity**

59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Historically, primary alcohols have been used as inhibitors of PLD hydrolysis activity. However, one must state that it is not a proper inhibitory effect as the use of alcohol derives PLD activity from hydrolysis *stricto sensu* toward the transphosphatidylation activity. In fact, alcohols can be used as secondary substrate and therefore PLD can transfer the phosphatidyl group to a primary alcohol to generate new phospholipids. The use of butan-1-ol is quite common to follow mammalian PLD activity by detecting the production of phosphatidylbutanol. We consequently tried to define which primary alcohol would be the best to avoid PLD-generated PA by testing the primary alcohols from methanol to hexan-1-ol (Fig. 4).

As expected, methanol and ethanol were not efficient secondary substrates of VuPLD. For short-chain alcohol the best candidate was butan-1-ol with an IC₅₀ of 150 mM compared to propan-1-ol which was more than twice less efficient with 320 mM. Pentan-1-ol had an IC₅₀ of 60 mM which is the best among all the alcohols tested. Finally, hexan-1-ol seemed not to be a proper secondary substrate as the activity quickly falls at 20 mM probably indicating enzyme denaturation (Fig. 4A). All secondary alcohols tested appeared not to be proper substrates as they displayed high IC₅₀, 160 mM for pentan-2-ol, 360 mM for butan-2-ol and propan-2-ol had no effect on PLD activity (Fig. 4A).

However, these values are based on the fluorimetric detection of PA (see Material & methods section). A more convenient way to detect phosphatidylalcohols issued from the PLD-catalyzed transphosphatidylation reaction is to use TLC for phospholipids separation. As can be seen from Fig. 4B, from all alcohols tested, the transphosphatidylation products were qualitatively detected only with methanol, ethanol, propan-1-ol and butan-1-ol.

281

Effect of known mammalian PLD inhibitors on VuPLD

1
2
3
4 283 Halopemide was shown previously to inhibit both human PLD1 and PLD2 isoforms, VU0359595
5
6 284 targets the PLD1 isoform with an IC_{50} calculated to be 3.7 nM, VU0285655 targets the PLD2
7
8
9 285 isoform with an IC_{50} calculated to be 635 nM. When tested on VuPLD, no inhibition could be
10
11 286 observed up to 10 μ M (Fig. 5). The PLD activity was reduced down to 50 % when 100 μ M of each
12
13
14 287 these three inhibitors were used independently.

15
16 288

19 289 **Test of alleged plant PLD inhibitors**

20
21 290 Amongst the molecules reported in literature to inhibit plant PLD activity we tested both
22
23 291 ciclosporin A and resveratrol on VuPLD using our fluorescence-based assay. In our hand, neither
24
25
26 292 ciclosporin A (Fig. 6A) nor resveratrol (Fig. 6B) were able to significantly affect VuPLD activity.
27
28
29 293 A slight 20% PLD inhibition could be observed using 2 mM of resveratrol, which is far from the
30
31 294 value reported by others where more than 90% of inhibition was measured with 130 μ M of
32
33 295 resveratrol [39]. As far as ciclosporin A is concerned, no inhibition was detected until 100 μ M
34
35
36 296 even if a quasi-total inhibition was previously reported for the PLD from cabbage using 5 μ M of
37
38 297 this compound [24,39].

39
40
41 298 Similarly, neither NAE (Fig. 6C) nor hexanal (Fig. 6D) which have been demonstrated as efficient
42
43 299 inhibitors of castor bean [21] or raspberry [29] PLDs, respectively, displayed any inhibitory effect
44
45 300 on VuPLD.

46
47 301

50 302 **Screening of new PLD inhibitors**

51
52
53 303 Because of its sensitivity and ease of use, we aimed to validate the 8HQ-based assay suitability for
54
55 304 HTS of novel PLD inhibitors as a start. In order to do so, we calculated the Z' -factor, a statistical
56
57
58 305 parameter used to evaluate the quality and performance of an assay. This parameter is reflective
59
60
61
62
63
64
65

1
2
3
4 306 of both the assay signal dynamic range and the data variation associated with signal measurements
5
6 307 [50]. In light of former results showing sodium vanadate as the most potent inhibitor of VuPLD
7
8
9 308 (Fig. 2B), we consequently decided to use it as a reference and a positive control for the screening.
10
11 309 Briefly, confronting the variability of 30 values of PLD activity measured with either 100 μ M of
12
13
14 310 vanadate or just 3 % DMSO, we obtained a calculated value of 0.803 for the Z' -score (Fig. 7A).
15
16 311 This value being comprised in the in the interval between 0.5 and 1.0 thus qualifies the 8HQ-based
17
18
19 312 test as “excellent” [50] for HTS of PLD inhibitors.
20
21 313 Moreover, as most of chemical library are DMSO solubilized, DMSO was shown only to have a
22
23
24 314 slight effect on PLD activity up to 5 % DMSO final concentration, but no statistical difference
25
26 315 compared from the control was observed when 5% DMSO final concentration was used (data not
27
28
29 316 shown).
30
31 317 We built a six-step protocol for the HTS of a PLD inhibitor (Fig. 7B). We consequently performed
32
33 318 a small-scale screening on the chemical library of our institute that is composed of 3,120
34
35
36 319 compounds from diverse origins. These molecules include quinolones, indoles, indolinones,
37
38 320 benzofurans, coumarins and flavonoids from chemistry and pharmaceutical laboratories.
39
40
41 321 A hit was considered every molecule capable to activate or inhibit PLD activity *in vitro* by more
42
43 322 than 50%. Out of 3,120 initial compounds, 227 molecules were found to reduce the PLD activity
44
45
46 323 and 53 to activate this activity (the others were found to have little or no effect at all). Because
47
48 324 some of the hits could be fluorescent molecules, or molecules with putative screen effect on the
49
50
51 325 8HQ probe, or molecules that could chelate the calcium ions for example, *id est* molecules that
52
53 326 could interfere with the fluorescent assay, we screened the putative 277 inhibitory molecules using
54
55
56 327 chromogenic assay which quantifies the free choline released by the PLD. After this step, 11
57
58 328 molecules were retained then further checked for their absence of effect on a PA standard curve
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

329 fluorescence. Finally, the 2 remaining compounds were tested on crude extract of *Pichia pastoris*
330 expressing VuPLD to avoid unspecific protein interaction. The last hit was confirmed against other
331 HKD-type PLDs like cabbage PLD [51] or peanut PLD [52]. Then, it was excluded after the last
332 step of this protocol as *de novo* synthesis of the molecule showed no effect at all and as several
333 contaminants were present in the initial batch of molecule used to make up the library.

335 Discussion

336 Using two different *ab initio* modelization programs SAXS data indicate that VuPLD is a globular
337 enzyme with elongated tops, thus the enzyme seems overall symmetrical with an ovoid shape.
338 These models do not allow the positioning of the regulatory C2 domain. The determination of the
339 overall shape of VuPLD is in accordance with the results obtained with cabbage PLD α 2 in 2012
340 by others [53]. Both models of nearly the same length (809 residues for VuPLD compare to 812
341 for cabbage PLD α 2) share 81.3% amino acid sequence identity and have a similar gyration radius,
342 but the D_{max} is lower in cabbage PLD (2.81 +/- 0.05 nm (this study) vs 2.89 +/- 0.01 nm [53])
343 indicating a slightly smaller enzyme. Also, the porod volume of VuPLD is more important
344 indicating a larger enzyme of 140.8 nm³ vs 126.7 nm³ for cabbage PLD [53]. These very slight
345 differences for such similar enzymes in terms of sequence might be explained by the fact that, even
346 if they are purified the same way, they are not recombinantly expressed in the same organism.
347 Specifically, cabbage PLD was expressed in *Escherichia coli* BL21 (DE3), contrary to VuPLD
348 that was expressed in yeast *Pichia pastoris* X33 cells which are known to glycosylate proteins.
349 The structure of PLD α 1 from Arabidopsis that was recently solved [12] positioned the C2 domain
350 as a small protuberance interacting with the second HKD domain which is, however, not in
351 accordance with the ovoid shape with extended extremities of the VuPLD obtained in this work.

1
2
3
4 352 Nevertheless, the SAXS model of VuPLD (this work) and of PLD from cabbage [53], or crystal
5
6 353 structure of *Arabidopsis* PLD [12] are very close (Fig. S2B) and interchangeable, which makes of
7
8
9 354 VuPLD another PLD model, and specially a convenient tool to study PLD inhibition as this latest
10
11
12 355 protein could be recombinantly expressed and easily purified in one step [54].
13
14 356 To prove that VuPLD is a relevant model to study plant PLD inhibition, we started with the use of
15
16 357 substrate analogs, in particular, analogs of the phosphate head of phospholipid. Therefore, we
17
18
19 358 tested phosphate salt at first sight and structurally close chemicals like vanadate or tungstate. These
20
21 359 salts are known to modulate PLD activity and we show here that vanadate is the best of them with
22
23
24 360 the finest IC₅₀. Consequently, we tested vanadate derivatives that differ by their hydrophobic
25
26 361 properties. Indeed, vanadate and then vanadate oxytriethoxide, vanadate oxytripropoxide,
27
28
29 362 vanadate oxytriisopropoxide and tris(triphenylsiloxy) vanadium oxide are increasing hydrophobic
30
31 363 molecules. Among them, vanadate oxytriethoxide that only display ethane moieties around
32
33
34 364 vanadate salt is the best inhibitor. These results may suggest that the binding site of vanadate on
35
36 365 PLD was not influenced by the alkyl chain length and/or steric environment and has enough room
37
38 366 to allow the fixation of the different derivatives tested.
39
40
41 367 Another way to inhibit the hydrolase activity of PLDs is the use of alcohols, keeping in mind that
42
43 368 alcohols are not proper inhibitors but are just another secondary substrate competing with water to
44
45
46 369 induce a transphosphatidylolation reaction. Here, we show that short-chained alcohols like methanol,
47
48 370 ethanol and to a lesser extent propan-1-ol do not affect the hydrolysis activity. Secondary alcohols
49
50
51 371 are also not substrates. The commonly used butan-1-ol is definitely the best primary alcohol
52
53 372 susceptible to generate a phosphatidylbutanol at a weak concentration. With other alcohols, no
54
55 373 transphosphatidylolation products were visible on TLC plates, meaning that they are not proper
56
57
58
59
60
61
62
63
64
65

1
2
3
4 374 secondary substrates of PLD. But one has to notice the fact that they indeed reduce PLD activity.
5
6 375 As stated above for hexan-1-ol, propan-2-ol and pentan-1-ol probably induce enzyme denaturation.
7
8
9 376 Human PLD inhibitors do not show any inhibitory effect on plant PLD, even if these PLD [11]
10
11 377 [12] share a common and superposable active site (Fig. S5). These results ask questions about the
12
13
14 378 mechanism by which they inhibit human PLDs. We demonstrated recently that all PLDs share
15
16 379 common sequence motifs [9]. Among them a pocket constituted by two HKD motifs form the
17
18
19 380 catalytic site. But also, several similar sequences are found in all eukaryotic PLDs. Indeed, PLDs
20
21 381 are mostly differing by their regulatory motifs [9]. For example, human PLD bears a PX and PH
22
23
24 382 domain [2] whereas plant PLD bears a C2-domain responsible for the fixation of Ca²⁺ ions, a
25
26 383 mandatory activator of plant PLD [55]. However, the catalytic sites of *Arabidopsis* PLD [12] and
27
28
29 384 human PLD1 [13] share a common feature as globular fold with a funnel-shaped cavity leading to
30
31 385 the active site. Considering the localized relative sequence similarity between human PLD and
32
33 386 plant PLD, and considering that if human PLD inhibitors were targeting the catalytic site they
34
35
36 387 would have a similar effect on plant PLD, it is possible that human PLD inhibitors are targeting
37
38 388 (i) aminoacid sequences that do not exist in plant PLD or (ii) the regulatory domains and therefore
39
40
41 389 do not inhibit the catalytic activity of the enzyme but its capacity to bind its partners and cofactors.
42
43 390 The structure of the catalytic domains of human PLD1 and PLD2 that was recently published [11]
44
45
46 391 would surely help to explain the selectivity of several molecules thought or used as generic
47
48 392 inhibitors on plant and/or mammalian PLDs. As it was not been possible to solve the structure of
49
50
51 393 PLD α 1 with FIPI [12], this compound is assumed to localized in the substrate-binding pocket of
52
53 394 the enzyme. However, our direct and continuous fluorescence based assay failed to display an
54
55 395 inhibitory effect on VuPLD, as no inhibition could be observed up to 100 μ M (Fig. 5) and only an
56
57
58 396 apparent 29% reduction of activity was observed when 1 mM of FIPI was used (data not shown),
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

397 noting that such a high concentration would probably interfere with the fluorescence of the 8HQ
398 probe. These differences of FIPI effects could be explained by the nature of the plant PLDs used,
399 as some residues are dissimilar in this pocket among the plant PLDs, or by the nature of the test,
400 the activity being assayed in a direct and continuous manner in this work.

401 At the exception of vanadate that is still the more potent plant PLD inhibitor to our knowledge,
402 almost of all plant PLD inhibitors described in the literature failed to inhibit VuPLD, consequently
403 there is still a need for a specific plant PLD inhibitor or a generalist eukaryotic PLD inhibitors
404 acting both on plant and mammalian PLDs. Looking closely, it appears that the use of a purified
405 enzyme in this study is the major difference compared to all other papers published on this topic.
406 In this regard, it is possible that these compounds interfere with others enzymes used in the cascade
407 of choline quantification. On the other hand, in our case the use of an assay to directly measure the
408 PA released directly from PLD-catalysed hydrolysis, constitutes a significant advantage to
409 quantify PLD activity [49].

410 Indeed, this 8HQ-based assay was statistically validated with a Z' score of 0.803 indicating an
411 excellent test for HTS and demonstrating that vanadate should be used as a positive control for
412 forthcoming screenings. We consequently screened as a proof of concept a chemical library of
413 ~3,120 compounds. After six runs of enzymatic tests combining successively the advantages of
414 both fluorescent and chromogenic assays to avoid false positives, no hit could unfortunately be
415 retained as the unique potential candidate appeared to be a contaminant. In the future, combining
416 the two PLD enzymatic assays with their specific strengths and weaknesses, their distinct and
417 alternative properties, larger libraries could be screened to identify and validate new PLD
418 inhibitors.

419

1
2
3
4 420 In conclusion, this work highlights (i) the necessity for a novel universal inhibitor potentially
5
6 421 targeting the catalytic site of PLDs, (ii) the potential of a purified plant PLD to screen PLD
7
8
9 422 inhibitors and (iii) the need to screen larger chemical libraries to find new inhibitory molecules.
10

11 423

12 424

13
14 425

15
16 426 Author Information

17
18
19 427 **Corresponding Author**

20
21 428 *To whom correspondence should be addressed

22
23
24 429 Tel: +33 4 27 46 57 31

25
26
27 430 E-mail address: alexandre.noiriel@univ-lyon1.fr

28
29
30 431

31
32
33 432 **Author Contributions**

34
35
36 433 YA and MVC performed SAXS and modeling experiments. AC and FP contributed to compound
37
38 434 design and synthesis. YA, KB, HA, MA, RR performed biochemical assays. LB, SM, AA and AN
39
40 435 devised the project and the main conceptual ideas. AN was in charge of funding acquisition. All
41
42 436 authors contributed to data interpretation. The manuscript was written from contributions by all
43
44 437 the authors. All the authors have given approval to the final version of the manuscript.
45
46

47
48
49 438

50
51
52 439 **Competing interests**

53
54
55 440 The authors have no competing interests to declare.
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

441

442

Acknowledgments

This work was funded by La Ligue contre le Cancer, comité de Haute-Savoie (France) and La Ligue contre le Cancer, comité de Saône-et-Loire (France). Houda Abla and Karim Bessaa were both supported by a grant by La Ligue contre le Cancer, comité de Haute-Savoie (France), Meryem Aydin was supported by La Ligue contre le Cancer, comité de Saône-et-Loire (France). Yani Arhab was supported by an MENRT grant from the French Ministry of Science and Education.

449

Abbreviations

8-Hydroxyquinoline (8HQ), high-throughput screening (HTS), lysophosphatidylethanolamine (LPE), N-acylethanolamines (NAE), N-acylphosphatidylethanolamine (NAPE), phosphatidic acid (PA), phospholipase D (PLD), palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC).

454

References

[1] S.M. Hammond, Y.M. Altshuller, T.C. Sung, S.A. Rudge, K. Rose, J. Engebrecht, A.J. Morris, M.A. Frohman, Human ADP-ribosylation factor-activated phosphatidylcholine-specific phospholipase D defines a new and highly conserved gene family, *J. Biol. Chem.* 270 (1995) 29640–29643.

[2] P.E. Selvy, R.R. Lavieri, C.W. Lindsley, H.A. Brown, Phospholipase D: enzymology, functionality, and chemical modulation, *Chem. Rev.* 111 (2011) 6064–6119. <https://doi.org/10.1021/cr200296t>.

- 1
2
3
4 463 [3] A. Marchler-Bauer, Y. Bo, L. Han, J. He, C.J. Lanczycki, S. Lu, F. Chitsaz, M.K. Derbyshire,
5 464 R.C. Geer, N.R. Gonzales, M. Gwadz, D.I. Hurwitz, F. Lu, G.H. Marchler, J.S. Song, N.
6 465 Thanki, Z. Wang, R.A. Yamashita, D. Zhang, C. Zheng, L.Y. Geer, S.H. Bryant,
7 466 CDD/SPARCLE: functional classification of proteins via subfamily domain architectures,
8 467 *Nucleic Acids Res.* 45 (2017) D200–D203. <https://doi.org/10.1093/nar/gkw1129>.
9
10 468 [4] S. Imamura, Y. Horiuti, Purification of *Streptomyces chromofuscus* phospholipase D by
11 469 hydrophobic affinity chromatography on palmitoyl cellulose, *J. Biochem. (Tokyo)*. 85 (1979)
12 470 79–95.
13
14 471 [5] S. Lee, K.R. Lynch, Brown recluse spider (*Loxosceles reclusa*) venom phospholipase D
15 472 (PLD) generates lysophosphatidic acid (LPA), *Biochem. J.* 391 (2005) 317–323.
16 473 <https://doi.org/10.1042/BJ20050043>.
17
18 474 [6] S.A. Rudge, J. Engebrecht, Regulation and function of PLDs in yeast, *Biochim. Biophys.*
19 475 *Acta.* 1439 (1999) 167–174.
20 476 [7] K. Rose, S.A. Rudge, M.A. Frohman, A.J. Morris, J. Engebrecht, Phospholipase D signaling
21 477 is essential for meiosis, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 12151–12155.
22
23 478 [8] I. Leiros, F. Secundo, C. Zambonelli, S. Servi, E. Hough, The first crystal structure of a
24 479 phospholipase D, *Struct. Lond. Engl.* 1993. 8 (2000) 655–667.
25 480 [9] Y. Arhab, A. Abousalham, A. Noiriél, Plant phospholipase D mining unravels new conserved
26 481 residues important for catalytic activity, *Biochim. Biophys. Acta Mol. Cell Biol. Lipids.* 1864
27 482 (2019) 688–703. <https://doi.org/10.1016/j.bbali.2019.01.008>.
28
29 483 [10] C.P. Ponting, I.D. Kerr, A novel family of phospholipase D homologues that includes
30 484 phospholipid synthases and putative endonucleases: identification of duplicated repeats and
31 485 potential active site residues, *Protein Sci. Publ. Protein Soc.* 5 (1996) 914–922.
32 486 <https://doi.org/10.1002/pro.5560050513>.
33
34 487 [11] C.M. Metrick, E.A. Peterson, J.C. Santoro, I.J. Enyedy, P. Murugan, T. Chen, K. Michelsen,
35 488 M. Cullivan, K.A. Spilker, P.R. Kumar, T.L. May-Dracka, J.V. Chodaparambil, Human PLD
36 489 structures enable drug design and characterization of isoenzyme selectivity, *Nat. Chem. Biol.*
37 490 (2020). <https://doi.org/10.1038/s41589-019-0458-4>.
38
39 491 [12] J. Li, F. Yu, H. Guo, R. Xiong, W. Zhang, F. He, M. Zhang, P. Zhang, Crystal structure of
40 492 plant PLD α 1 reveals catalytic and regulatory mechanisms of eukaryotic phospholipase D,
41 493 *Cell Res.* (2019). <https://doi.org/10.1038/s41422-019-0244-6>.
42 494 [13] F.Z. Bowling, C.M. Salazar, J.A. Bell, T.S. Huq, M.A. Frohman, M.V. Airola, Crystal
43 495 structure of human PLD1 provides insight into activation by PI(4,5)P₂ and RhoA, *Nat. Chem.*
44 496 *Biol.* 16 (2020) 400–407. <https://doi.org/10.1038/s41589-020-0499-8>.
45
46 497 [14] M.I. McDermott, Y. Wang, M.J.O. Wakelam, V.A. Bankaitis, Mammalian phospholipase D:
47 498 Function, and therapeutics, *Prog. Lipid Res.* (2019) 101018.
48 499 <https://doi.org/10.1016/j.plipres.2019.101018>.
49
50 500 [15] A. Kulkarni, P. Quang, V. Curry, R. Keyes, W. Zhou, H. Cho, J. Baffoe, B. Török, K.
51 501 Stieglitz, 1,3-disubstituted-4-aminopyrazolo [3, 4-d] pyrimidines, a new class of potent
52 502 inhibitors for phospholipase D, *Chem. Biol. Drug Des.* 84 (2014) 270–281.
53 503 <https://doi.org/10.1111/cbdd.12319>.
54 504 [16] H. Djakpa, A. Kulkarni, S. Barrows-Murphy, G. Miller, W. Zhou, H. Cho, B. Török, K.
55 505 Stieglitz, Identifying New Drug Targets for Potent Phospholipase D Inhibitors: Combining
56 506 Sequence Alignment, Molecular Docking, and Enzyme Activity/Binding Assays, *Chem.*
57 507 *Biol. Drug Des.* 87 (2016) 714–729. <https://doi.org/10.1111/cbdd.12705>.

- 1
2
3
4 508 [17] D.K. Behera, P.M. Behera, L. Acharya, A. Dixit, Pharmacophore modelling, virtual
5 509 screening and molecular docking studies on PLD1 inhibitors, SAR QSAR Environ. Res. 28
6 510 (2017) 991–1009. <https://doi.org/10.1080/1062936X.2017.1393774>.
- 8 511 [18] L. Monovich, B. Mugrage, E. Quadros, K. Toscano, R. Tommasi, S. LaVoie, E. Liu, Z. Du,
9 512 D. LaSala, W. Boyar, P. Steed, Optimization of halopemide for phospholipase D2 inhibition,
10 513 Bioorg. Med. Chem. Lett. 17 (2007) 2310–2311. <https://doi.org/10.1016/j.bmcl.2007.01.059>.
- 12 514 [19] Y. Ben Ali, F. Carriere, A. Abousalham, High-level constitutive expression in *Pichia pastoris*
13 515 and one-step purification of phospholipase D from cowpea (*Vigna unguiculata* L. Walp),
14 516 Protein Expr Purif. 51 (2007) 162–9. <https://doi.org/10.1016/j.pep.2006.07.018>.
- 15 517 [20] Z. Hussain, T. Uyama, K. Tsuboi, N. Ueda, Mammalian enzymes responsible for the
16 518 biosynthesis of N-acylethanolamines, Biochim. Biophys. Acta. 1862 (2017) 1546–1561.
17 519 <https://doi.org/10.1016/j.bbali.2017.08.006>.
- 19 520 [21] S.L. Austin-Brown, K.D. Chapman, Inhibition of phospholipase D alpha by N-
20 521 acylethanolamines, Plant Physiol. 129 (2002) 1892–1898.
21 522 <https://doi.org/10.1104/pp.001974>.
- 22 523 [22] H. Chen, X. Yu, X. Zhang, L. Yang, X. Huang, J. Zhang, H.W. Pritchard, W. Li,
23 524 Phospholipase D α 1-mediated phosphatidic acid change is a key determinant of desiccation-
24 525 induced viability loss in seeds, Plant Cell Environ. 41 (2018) 50–63.
25 526 <https://doi.org/10.1111/pce.12925>.
- 27 527 [23] S.B. Ryu, B.H. Karlsson, M. Ozgen, J.P. Palta, Inhibition of phospholipase D by
28 528 lysophosphatidylethanolamine, a lipid-derived senescence retardant, Proc. Natl. Acad. Sci.
29 529 U. S. A. 94 (1997) 12717–12721.
- 31 530 [24] M. Madesh, K.A. Balasubramanian, Cyclosporin A inhibits oxidant and calcium stimulated
32 531 phospholipase D activity in the rat intestinal mitochondria, Biochim. Biophys. Acta. 1389
33 532 (1998) 206–212.
- 34 533 [25] N. Dittrich, G. Nössner, B. Kutscher, R. Ulbrich-Hofmann, Alkylphosphate esters as
35 534 inhibitors of phospholipase D, J. Enzym. Inhib. 11 (1996) 67–75.
- 37 535 [26] Y. Ohno, M. Okazaki, I. Hara, Studies on the activity of phospholipase D with alkyl
38 536 phosphorylcholines, Biochem. Cell Biol. Biochim. Biol. Cell. 68 (1990) 376–379.
- 39 537 [27] G. Paliyath, R.G. Pinhero, R.Y. Yada, D.P. Murr, Effect of processing conditions on
40 538 phospholipase D activity of corn kernel subcellular fractions, J. Agric. Food Chem. 47 (1999)
41 539 2579–2588.
- 43 540 [28] K. Tiwari, G. Paliyath, Microarray analysis of ripening-regulated gene expression and its
44 541 modulation by 1-MCP and hexanal, Plant Physiol. Biochem. PPB. 49 (2011) 329–340.
45 542 <https://doi.org/10.1016/j.plaphy.2011.01.007>.
- 47 543 [29] W.E. Kayal, G. Paliyath, J.A. Sullivan, J. Subramanian, Phospholipase D inhibition by
48 544 hexanal is associated with calcium signal transduction events in raspberry, Hort. Res. 4
49 545 (2017) 17042. <https://doi.org/10.1038/hortres.2017.42>.
- 50 546 [30] B. Bhushan, S. Kumar, M.K. Mahawar, K. Jalgaonkar, A.S. Dukare, B. Bibwe, V.S. Meena,
51 547 N. Negi, R.K. Narwal, A. Pal, Nullifying phosphatidic acid effect and controlling
52 548 phospholipase D associated browning in litchi pericarp through combinatorial application of
53 549 hexanal and inositol, Sci. Rep. 9 (2019) 2402. <https://doi.org/10.1038/s41598-019-38694-5>.
- 54 550 [31] D. Dhakshinamoorthy, S. Sundaresan, A. Iyadurai, K.S. Subramanian, G.J. Janavi, G.
55 551 Paliyath, J. Subramanian, Hexanal Vapor Induced Resistance against Major Postharvest
56 552 Pathogens of Banana (*Musa acuminata* L.), Plant Pathol. J. 36 (2020) 133–147.
57 553 <https://doi.org/10.5423/PPJ.OA.03.2019.0072>.
- 60
61
62
63
64
65

- 1
2
3
4 554 [32] A. Kaur, K. Neelam, K. Kaur, A. Kitazumi, B.G. de Los Reyes, K. Singh, Novel allelic
5 555 variation in the Phospholipase D alpha1 gene (OsPLD α 1) of wild *Oryza* species implies to
6 556 its low expression in rice bran, *Sci. Rep.* 10 (2020) 6571. [https://doi.org/10.1038/s41598-](https://doi.org/10.1038/s41598-020-62649-w)
7 557 [020-62649-w](https://doi.org/10.1038/s41598-020-62649-w).
- 9 558 [33] S. Lukowski, J.P. Mira, A. Zachowski, B. Geny, Fodrin inhibits phospholipases A2, C, and
10 559 D by decreasing polyphosphoinositide cell content, *Biochem. Biophys. Res. Commun.* 248
11 560 (1998) 278–284. <https://doi.org/10.1006/bbrc.1998.8942>.
- 13 561 [34] J.A. Lewis, S.A. Scott, R. Lavieri, J.R. Buck, P.E. Selvy, S.L. Stoops, M.D. Armstrong, H.A.
14 562 Brown, C.W. Lindsley, Design and synthesis of isoform-selective phospholipase D (PLD)
15 563 inhibitors. Part I: Impact of alternative halogenated privileged structures for PLD1
16 564 specificity, *Bioorg. Med. Chem. Lett.* 19 (2009) 1916–1920.
17 565 <https://doi.org/10.1016/j.bmcl.2009.02.057>.
- 19 566 [35] S.A. Scott, P.E. Selvy, J.R. Buck, H.P. Cho, T.L. Criswell, A.L. Thomas, M.D. Armstrong,
20 567 C.L. Arteaga, C.W. Lindsley, H.A. Brown, Design of isoform-selective phospholipase D
21 568 inhibitors that modulate cancer cell invasiveness, *Nat. Chem. Biol.* 5 (2009) 108–117.
22 569 <https://doi.org/10.1038/nchembio.140>.
- 24 570 [36] W. Su, O. Yeku, S. Olepu, A. Genna, J.-S. Park, H. Ren, G. Du, M.H. Gelb, A.J. Morris,
25 571 M.A. Frohman, 5-Fluoro-2-indolyl des-chlorohalopemide (FIPI), a phospholipase D
26 572 pharmacological inhibitor that alters cell spreading and inhibits chemotaxis, *Mol. Pharmacol.*
27 573 75 (2009) 437–446. <https://doi.org/10.1124/mol.108.053298>.
- 29 574 [37] R.R. Lavieri, S.A. Scott, P.E. Selvy, K. Kim, S. Jadhav, R.D. Morrison, J.S. Daniels, H.A.
30 575 Brown, C.W. Lindsley, Design, synthesis, and biological evaluation of halogenated N-(2-(4-
31 576 oxo-1-phenyl-1,3,8-triazaspiro[4.5]decan-8-yl)ethyl)benzamides: discovery of an isoform-
32 577 selective small molecule phospholipase D2 inhibitor, *J. Med. Chem.* 53 (2010) 6706–6719.
33 578 <https://doi.org/10.1021/jm100814g>.
- 35 579 [38] R. Ganesan, M. Mahankali, G. Alter, J. Gomez-Cambronero, Two sites of action for PLD2
36 580 inhibitors: The enzyme catalytic center and an allosteric, phosphoinositide binding pocket,
37 581 *Biochim. Biophys. Acta.* 1851 (2015) 261–272.
38 582 <https://doi.org/10.1016/j.bbali.2014.12.007>.
- 39 583 [39] S. Majd, E.C. Yusko, A.D. MacBriar, J. Yang, M. Mayer, Gramicidin pores report the activity
40 584 of membrane-active enzymes, *J. Am. Chem. Soc.* 131 (2009) 16119–16126.
41 585 <https://doi.org/10.1021/ja904072s>.
- 43 586 [40] J. Tou, C. Urbizo, Resveratrol inhibits the formation of phosphatidic acid and diglyceride in
44 587 chemotactic peptide- or phorbol ester-stimulated human neutrophils, *Cell. Signal.* 13 (2001)
45 588 191–197. [https://doi.org/10.1016/s0898-6568\(01\)00137-1](https://doi.org/10.1016/s0898-6568(01)00137-1).
- 47 589 [41] L. Li, N. Fleming, Aluminum fluoride inhibition of cabbage phospholipase D by a phosphate-
48 590 mimicking mechanism, *FEBS Lett.* 461 (1999) 1–5. [https://doi.org/10.1016/s0014-](https://doi.org/10.1016/s0014-5793(99)01414-3)
49 591 [5793\(99\)01414-3](https://doi.org/10.1016/s0014-5793(99)01414-3).
- 50 592 [42] M.M. Bradford, A rapid and sensitive method for the quantitation of microgram quantities of
51 593 protein utilizing the principle of protein-dye binding, *Anal Biochem.* 72 (1976) 248–54.
- 53 594 [43] D.I. Svergun, Restoring low resolution structure of biological macromolecules from solution
54 595 scattering using simulated annealing, *Biophys. J.* 76 (1999) 2879–2886.
55 596 [https://doi.org/10.1016/S0006-3495\(99\)77443-6](https://doi.org/10.1016/S0006-3495(99)77443-6).
- 57 597 [44] D.I. Svergun, M.V. Petoukhov, M.H. Koch, Determination of domain structure of proteins
58 598 from X-ray solution scattering, *Biophys. J.* 80 (2001) 2946–2953.
59 599 [https://doi.org/10.1016/S0006-3495\(01\)76260-1](https://doi.org/10.1016/S0006-3495(01)76260-1).

- 1
2
3
4 600 [45] V.V. Volkov, Svergun, D.I., Uniqueness of ab initio shape determination in small-angle
5 601 scattering, *J. Appl. Crystallogr.* (2003) 860–864.
6 602 [46] D. Schneidman-Duhovny, M. Hammel, J.A. Tainer, A. Sali, FoXS, FoXSDock and
7 603 MultiFoXS: Single-state and multi-state structural modeling of proteins and their complexes
8 604 based on SAXS profiles, *Nucleic Acids Res.* 44 (2016) W424–429.
9 605 <https://doi.org/10.1093/nar/gkw389>.
10 606 [47] V. Piiadov, E. Ares de Araújo, M. Oliveira Neto, A.F. Craievich, I. Polikarpov, SAXSMoW
11 607 2.0: Online calculator of the molecular weight of proteins in dilute solution from experimental
12 608 SAXS data measured on a relative scale, *Protein Sci. Publ. Protein Soc.* 28 (2019) 454–463.
13 609 <https://doi.org/10.1002/pro.3528>.
14 610 [48] Y. Arhab, R. Rahier, A. Noiriél, M.V. Cherrier, A. Abousalham, Expression and Purification
15 611 of Recombinant *Vigna unguiculata* Phospholipase D in *Pichia pastoris* for Structural Studies,
16 612 in: G. Sandoval (Ed.), *Lipases Phospholipases*, Springer New York, New York, NY, 2018:
17 613 pp. 191–201. https://doi.org/10.1007/978-1-4939-8672-9_10.
18 614 [49] R. Rahier, A. Noiriél, A. Abousalham, Development of a Direct and Continuous
19 615 Phospholipase D Assay Based on the Chelation-Enhanced Fluorescence Property of 8-
20 616 Hydroxyquinoline, *Anal. Chem.* 88 (2016) 666–74.
21 617 <https://doi.org/10.1021/acs.analchem.5b02332>.
22 618 [50] J. Zhang, T. Chung, K. Oldenburg, A Simple Statistical Parameter for Use in Evaluation and
23 619 Validation of High Throughput Screening Assays, *J. Biomol. Screen.* 4 (1999) 67–73.
24 620 <https://doi.org/10.1177/108705719900400206>.
25 621 [51] H. Sato, T. Watanabe, Y. Sagane, Y. Nakazawa, K. Takano, Purification and Characterization
26 622 of Phospholipase D from Cabbage Leaves., *Food Sci. Technol. Res.* 6 (2000) 29–33.
27 623 <https://doi.org/10.3136/fstr.6.29>.
28 624 [52] Y. Nakazawa, H. Sato, M. Uchino, K. Takano, Purification, characterization and cloning of
29 625 phospholipase D from peanut seeds, *Protein J.* 25 (2006) 212–223.
30 626 <https://doi.org/10.1007/s10930-006-9004-9>.
31 627 [53] S. Stumpe, S. König, R. Ulbrich-Hofmann, Insights into the structure of plant α -type
32 628 phospholipase D: Structure of plant phospholipase D, *FEBS J.* 274 (2007) 2630–2640.
33 629 <https://doi.org/10.1111/j.1742-4658.2007.05798.x>.
34 630 [54] Arhab, Y, Rahier, R, Noiriél, A, Cherrier, MV, Abousalham, A, Expression and Purification
35 631 of Recombinant *Vigna unguiculata* Phospholipase D in *Pichia pastoris* for Structural Studies,
36 632 *Methods Mol. Biol. Press.* (2018) In press.
37 633 [55] K. Pappan, X. Wang, Molecular and biochemical properties and physiological roles of plant
38 634 phospholipase D, *Biochim. Biophys. Acta.* 1439 (1999) 151–166.
39 635
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Fig. 1: SAXS modelization of VuPLD.

A) Diffusion curve. B) Kratky curve. C) Distance distribution function $P(r)$ calculated from X-ray scattering data. D) *Ab initio* low-resolution structure model of VuPLD calculated from the SAXS pattern and the program DAMMIN. Model is shown in two different views obtained by 90° rotation around the y-axis.




Fig. 2: Inhibitory effect of different salts on purified VuPLD.

The PLD activity was measured with the fluorimetric assay in triplicates and with increasing concentrations of sodium vanadate (A), sodium tungstate (B) or sodium phosphate (C). D) PLD relative activity incubated with 0 μM (circle ●), 25 μM (square ■), 50 μM (triangle ▲) or 100 μM (cross *) of phosphate at different concentrations of vanadate. Relative PLD activities were calculated relatively to the enzyme incubated in the absence of effector set at 100 %. Values are the mean \pm SD obtained from three independent assays.




Fig. 3: Inhibitory effect of different vanadate derivatives on purified VuPLD.

The activity was measured with the chromogenic assay in triplicates with increasing concentrations of vanadate (circle ●), vanadate oxytriethoxide (square ■), vanadate oxytripropoxide (triangle ▲), vanadate oxytriisopropoxide (cross ×) or vanadate tris(triphenylsiloxy) (diamond ◆). Values are the mean \pm SD obtained from three independent chromogenic assays.





Fig. 4: Effect of different alcohols on purified VuPLD.

A) The activity was measured in triplicates with increasing concentrations of methanol (black dash—), ethanol (square■), propan-1-ol (white triangle Δ), propan-2-ol (black triangle ▲), butan-1-ol (white diamond ◇), butan-2-ol (black diamond ◆), pentan-1-ol (white circle ○), pentan-2-ol (black circle ●) and hexanol (cross *). Relative activities were calculated relatively to the enzyme incubated in the absence of effector (100%). Values are the mean ± SD obtained from three independent fluorimetric assays. B) TLC analysis of the VuPLD catalyzed transphosphatidylation reaction products obtained from (A). Lipids were extracted, developed, and revealed as described in the experimental section. POPA, phosphatidylbutanol (P-But) and phosphatidylethanol (P-Eth) were run as standard reaction products,




Fig. 5: Inhibitory effect of known inhibitors of human PLD on purified plant VuPLD.

The activity was measured in triplicates with increasing concentrations of halopemide (black), VU0359595 (dark grey), VU0285655 (light grey), FIPI (white). Relative activities are calculated relatively to the enzyme incubated with 0 μM of effector (100%). Values are the mean ± SD obtained from three independent fluorimetric assays. ** P<0.01, *** P<0.001 (vs untreated).




Fig. 6: Inhibitory effect of alleged inhibitors of plant PLD.

PLD described inhibitors, ciclosporin A (A), resveratrol (B), NAE C12:0 (C) and hexanal (D) were tested at different concentrations from 1 μ M to 1 mM. Relative activities are then calculated relatively to the enzyme incubated in the absence of effector (100%). Values are the mean \pm SD obtained from three independent fluorimetric assays. * $P < 0.05$, ** $P < 0.01$ (vs untreated).




Fig. 7: HTS screening of a new PLD inhibitor.

PLD activity was measured in the presence or absence of 100 μM of vanadate in order to establish the variation of the assay and the calculation of the Z' score. Multiple repeats ($n=30$) were realized with (circles) or without effector (squares). B) Seven-step workflow for the identification of novel PLD inhibitors.




Fig. S1: SAXS modelization of VuPLD.
Ab initio low-resolution structure model of VuPLD calculated from the SAXS pattern and the program GASBOR. Model is shown in two different views obtained by 90° rotation around the y-axis




Fig. S2: Analysis of SAXS data of VuPLD.

A) Comparison of the experimental scattering data (gray circles) with the calculated scattering profile from PLD structure.

B) Experimental SAXS curve of VuPLD (grey dots) compared to the theoretical SAXS curve of PLD α 1 (black line) from Arabidopsis (PDB code 6KZ9).




Fig. S3: Chemical structures of vanadate and its derivatives




Fig. S4: Inhibition of purified plant PLD by vanadate. The activity was measured in triplicates with increasing concentrations of vanadate. Values are the mean \pm SD obtained from three independent chromogenic assays.




Fig. S5: Comparison of plant and mammalian PLD catalytic pockets. Both HKD motifs are displayed in blue (human PLD, 6OHR) and red (plant PLD, 6KZ9) according to [9]. Histidyl and lysyl catalytic residues are represented (sticks) and annotated. Structure superposition was done using Pymol 2.3.2..

Author Statement

YA and MVC performed SAXS and modeling experiments. AC and FP contributed to compound design and synthesis. YA, KB, HA, MA, RR performed biochemical assays. LB, SM, AA and AN devised the project and the main conceptual ideas. AN was in charge of funding acquisition. All authors contributed to data interpretation. The manuscript was written from contributions by all the authors. All the authors have given approval to the final version of the manuscript.