



**HAL**  
open science

## **Impact of COVID-19 pandemic on mental health: An international study**

Andrew Gloster, Demetris Lamnisis, Jelena Lubenko, Giovambattista Presti, Valeria Squatrito, Marios Constantinou, Christiana Nicolaou, Savvas Papacostas, Gökçen Aydın, Yuen Yu Chong, et al.

► **To cite this version:**

Andrew Gloster, Demetris Lamnisis, Jelena Lubenko, Giovambattista Presti, Valeria Squatrito, et al.. Impact of COVID-19 pandemic on mental health: An international study. PLoS ONE, 2020, 15 (12), pp.e0244809. 10.1371/journal.pone.0244809 . hal-03177597

**HAL Id: hal-03177597**

<https://hal.univ-grenoble-alpes.fr/hal-03177597v1>

Submitted on 23 Mar 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## RESEARCH ARTICLE

## Impact of COVID-19 pandemic on mental health: An international study

Andrew T. Gloster<sup>1†\*</sup>, Demetris Lamnisos<sup>2</sup>, Jelena Lubenko<sup>3</sup>, Giovambattista Presti<sup>4</sup>, Valeria Squatrito<sup>4</sup>, Marios Constantinou<sup>5</sup>, Christiana Nicolaou<sup>6</sup>, Savvas Papacostas<sup>7</sup>, Gökçen Aydın<sup>8</sup>, Yuen Yu Chong<sup>9</sup>, Wai Tong Chien<sup>9</sup>, Ho Yu Cheng<sup>9</sup>, Francisco J. Ruiz<sup>10</sup>, Maria B. Garcia-Martin<sup>11</sup>, Diana P. Obando-Posada<sup>11</sup>, Miguel A. Segura-Vargas<sup>10</sup>, Vasilis S. Vasiliou<sup>12</sup>, Louise McHugh<sup>13</sup>, Stefan Höfer<sup>14</sup>, Adriana Baban<sup>15</sup>, David Dias Neto<sup>16</sup>, Ana Nunes da Silva<sup>17</sup>, Jean-Louis Monestès<sup>18</sup>, Javier Alvarez-Galvez<sup>19</sup>, Marisa Paez-Blarrina<sup>20</sup>, Francisco Montesinos<sup>21</sup>, Sonsoles Valdivia-Salas<sup>22</sup>, Dorottya Ori<sup>23</sup>, Bartosz Kleszcz<sup>24</sup>, Raimo Lappalainen<sup>25</sup>, Iva Ivanović<sup>26</sup>, David Gosar<sup>27</sup>, Frederick Dionne<sup>28</sup>, Rhonda M. Merwin<sup>29</sup>, Angelos P. Kassianos<sup>30‡</sup>, Maria Karekla<sup>30‡</sup>

1 Division of Clinical Psychology & Intervention Science, Department of Psychology, University of Basel, Basel, Switzerland, 2 Department of Health Sciences, European University Cyprus, Nicosia, Cyprus, 3 Psychological Laboratory, Faculty of Public Health and Social Welfare, Riga Stradiņš University, Riga, Latvia, 4 Kore University Behavioral Lab (KUBeLab), Faculty of Human and Social Sciences, Kore University of Enna, Enna, Italy, 5 Department of Social Sciences, School of Humanities and Social Sciences, University of Nicosia, Nicosia, Cyprus, 6 Department of Nursing, Cyprus University of Technology, Limassol, Cyprus, 7 Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus, 8 Department of Psychological Counseling and Guidance, Faculty of Education, Hasan Kalyoncu University, Gaziantep, Turkey, 9 The Nethersole School of Nursing, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, 10 Department of Psychology, Fundación Universitaria Konrad Lorenz, Bogotá, Colombia, 11 Faculty of Psychology, University of La Sabana, Chía, Colombia, 12 School of Applied Psychology, University College Cork, Cork, Ireland, 13 School of Psychology, University College Dublin, Dublin, Ireland, 14 Medical University Innsbruck, Innsbruck, Austria, 15 Department of Psychology, Babeş-Bolyai University (UBB), Cluj-Napoca, Romania, 16 Instituto Superior de Psicologia Aplicada (ISPA), Instituto Universitário; APPsyCI—Applied Psychology Research Center Capabilities & Inclusion, Lisboa, Portugal, 17 Faculdade de Psicologia, Alameda da Universidade, Universidade de Lisboa, Lisboa, Portugal, 18 LIP/PC2S, Université Grenoble Alpes, Grenoble, France, 19 Department of Biomedicine, Biotechnology and Public Health, University of Cadiz, Cadiz, Spain, 20 Instituto ACT, Madrid, Spain, 21 Department of Psychology, European University of Madrid, Madrid, Spain, 22 Department of Psychology and Sociology, University of Zaragoza, Zaragoza, Spain, 23 Vadaskert Child and Adolescent Psychiatric Hospital, Budapest, Hungary, 24 Private Practice, Poland, 25 Department of Psychology, University of Jyväskylä, Jyväskylä, Finland, 26 Clinic for Psychiatry, Clinical Center of Montenegro, Podgorica, Montenegro, 27 Ljubljana University Medical Centre, Ljubljana, Slovenia, 28 Département de Psychologie, Université du Québec à Trois-Rivières, Trois-Rivières, Canada, 29 Department of Psychiatry and Behavioral Science, Duke University, Durham, North Carolina, United States of America, 30 Department of Psychology, University of Cyprus, Nicosia, Cyprus

† ATG, MK and AK designed and implemented the study together. AK and MK should be considered joint senior authors.

\* [andrew.gloster@unibas.ch](mailto:andrew.gloster@unibas.ch)



## OPEN ACCESS

**Citation:** Gloster AT, Lamnisos D, Lubenko J, Presti G, Squatrito V, Constantinou M, et al. (2020) Impact of COVID-19 pandemic on mental health: An international study. PLoS ONE 15(12): e0244809. <https://doi.org/10.1371/journal.pone.0244809>

**Editor:** Joel Msafiri Francis, University of the Witwatersrand, SOUTH AFRICA

**Received:** October 3, 2020

**Accepted:** December 16, 2020

**Published:** December 31, 2020

**Copyright:** © 2020 Gloster et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its [Supporting Information](#) files.

**Funding:** This work was supported by grants from the Swiss National Science Foundation awarded to Andrew T. Gloster (PP00P1\_163716/1 & PP00P1\_190082). The funder provided support in the form of salaries for authors [ATG], but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of

## Abstract

## Background

The COVID-19 pandemic triggered vast governmental lockdowns. The impact of these lockdowns on mental health is inadequately understood. On the one hand such drastic changes in daily routines could be detrimental to mental health. On the other hand, it might not be experienced negatively, especially because the entire population was affected.

these authors are articulated in the 'author contributions' section. One of the authors is employed by a commercial affiliation: Private Prattice, Poland. This affiliation provided support in the form of salaries for authors [BK], but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** One of the authors is employed by a commercial affiliation: Private Prattice, Poland. This affiliation provided support in the form of salaries for author BK, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. This does not alter our adherence to PLOS ONE policies on sharing data and materials. No other authors have competing interests to declare.

## Methods

The aim of this study was to determine mental health outcomes during pandemic induced lockdowns and to examine known predictors of mental health outcomes. We therefore surveyed  $n = 9,565$  people from 78 countries and 18 languages. Outcomes assessed were stress, depression, affect, and wellbeing. Predictors included country, sociodemographic factors, lockdown characteristics, social factors, and psychological factors.

## Results

Results indicated that on average about 10% of the sample was languishing from low levels of mental health and about 50% had only moderate mental health. Importantly, three consistent predictors of mental health emerged: social support, education level, and psychologically flexible (vs. rigid) responding. Poorer outcomes were most strongly predicted by a worsening of finances and not having access to basic supplies.

## Conclusions

These results suggest that on whole, respondents were moderately mentally healthy at the time of a population-wide lockdown. The highest level of mental health difficulties were found in approximately 10% of the population. Findings suggest that public health initiatives should target people without social support and those whose finances worsen as a result of the lockdown. Interventions that promote psychological flexibility may mitigate the impact of the pandemic.

## Introduction

The COVID-19 global pandemic caused by the severe acute respiratory syndrome coronavirus 2 (SARS-COV-2) virus triggered governmentally mandated lockdowns, social distancing, quarantines and other measures in the interest of public health. The mandated lockdowns abruptly and dramatically altered people's daily routines, work, travel, and leisure activities to a degree unexperienced by most people living outside of war zones. Simultaneously, the highly contagious, yet invisible virus transformed previously neutral situations to perceived potentially dangerous ones: social interaction, touching one's face, going to a concert, shaking someone's hand, and even hugging grandparents. Given these changes and looming threat, increases in anxiety and depression can be expected [1]. Indeed, common psychological reactions to previous quarantines include post-traumatic symptoms, confusion, and anger [2], though these data stem from quarantines of specific regions or a subgroup of exposed people, such as medical professionals. It therefore remains an empirical question whether such patterns are consistent when entire populations across the globe are simultaneously affected.

For most people, it stands to reason that governmentally mandated lockdowns decrease their activity levels and the number of stimuli experienced compared to pre-lockdown levels. The impact of reducing activities, stimuli and routines on the population is unknown, but various analogue situations can be used to make predictions, like death of a spouse [3]; hearing loss [4]; job loss [5]; long duration expeditions [6]; poor acculturation [7]; and even ageing when combined with loneliness [8]. Each of these situations is associated with increases in psychological distress. This reduction of stimulations may lead to boredom and reductions in reinforcement, which has been associated with depression [9]. The sum total of these

literatures, and some evidence from country specific studies on COVID-19 suggests that for some people, the mental distress in the form of stress, depression, and negative affect are likely reactions to the lockdown; therefore, people's wellbeing is likely to suffer. Indeed, increased loneliness, social isolation, and living alone are associated with increased mortality [10]—the exact effect that mandated lockdown and social distancing rules aimed to counteract.

Alternately, the planned slowing down of daily routines can be beneficial. For example, vacations and weekends are highly sought-after—if not always achieved—periods of relaxation and stress reduction [11]. Likewise, some religious and spiritual traditions encourage simplicity, mindfulness, and solitude with the goal of increasing wellbeing [12]. It is therefore conceivable that for some people the lockdown could offer a reprieve from daily hassles and stress and even lead to increases in wellbeing. It is therefore equally important to identify protective factors that can buffer against the negative effects of the lockdown.

Although nearly all people around the globe have been subject to some form of lockdown measures to contain the COVID-19 response, variations exist with respect to how each person is confined, even within a single country. For instance, during the COVID-19 pandemic some people were allowed to go to work, whereas others were required to work exclusively from home. For various reasons, some people had difficulty obtaining some basic supplies. Further, some were thrust into the situation of taking care of others (e.g., children, due to closing of schools). Finally, some people lost income as a result of the lockdown, and this is a known risk-factor for poor mental health [13, 14]. Finally, a lockdown may be experienced differently the longer it continues and potentially when in confined spaces [2]. All of these lockdown-specific features may have an impact on one's mental health, but to date it remains inadequately explored.

As the risk of the pandemic continues, it is important to understand to what degree the virus-induced uncertainty and the lockdown-induced changes in daily routines impact stress, depression, affect, and wellbeing. Towards this end, it is important to identify factors that can mitigate potential negative psychological effects of pandemics and lockdowns. Various social and psychological factors have been identified in other contexts that may also help build resilience in large-scale pandemics such as COVID-19. On the social level, one such candidate is social support, which has repeatedly been found to positively impact mental health and wellbeing [15–18]. Another social factor is the family climate and family functioning, which clearly impacts people's mental health [19, 20]. Psychological factors such as mindfulness and psychologically flexible response styles (as opposed to rigid and avoidant response styles) are behavioral repertoires that have previously been shown to buffer the impact of stress and facilitate wellbeing [21–24].

Given the scope of the COVID-19 pandemic, it is crucial to better understand how a pandemic and associated lockdowns impact on mental health. Thus, the aim of this study was to determine mental health outcomes and to examine known predictors of outcomes to identify psychological processes and contextual factors that can be used in developing public health interventions. It can be assumed, but remains untested, that those with risks in social-demographic factors, living conditions, social factors and psychological factors have more severe reactions to the lockdown. We therefore tested whether outcomes of stress, depression, affect, and wellbeing were predicted by country of residence, social demographic characteristics, COVID-19 lockdown related predictors, social predictors, and psychological predictors.

## Methods

### Participants

The inclusion criteria were  $\geq 18$  years of age and ability to read one of the 18 languages (English, Greek, German, French, Spanish, Turkish, Dutch, Latvian, Italian, Portuguese,

Finnish, Slovenian, Polish, Romanian, Hong Kong, Hungarian, Montenegrin, & Persian.). There were no exclusion criteria. People from all countries were eligible to participate.

## Procedure

Ethics approval was obtained from the Cyprus National Bioethics Committee (ref.: EEBK EΠ 2020.01.60) followed by site approvals from different research teams involved in data collection. All participants provided written informed consent prior to completing the survey (computer-based, e.g., by clicking “yes”).

A population based cross-sectional study was conducted in order to explore how people across the world reacted to the COVID-19. The anonymous online survey was distributed using a range of methods. Universities emailed the online survey to students and academic staff and also posted the survey link to their websites. In addition, and in order to broaden the sample to older age groups and to those with different socio-demographic characteristics, the survey was disseminated in local press (e.g., newspapers, newsletters, radio stations), in social media (e.g., Facebook, Twitter, etc.), in professional networks, local hospitals and health centers and professional groups' email lists (e.g., medical doctors, teachers, engineers, psychologists, government workers), and to social institutions in the countries (e.g., churches, schools, cities/townships, clubs, etc.).

Data were collected for two months between 07th April and 07th June 2020. The majority of countries where data were collected had declared a state of emergency for COVID-19 during this time.

## Measures

Well validated and established measures were used to assess constructs. When measures did not already exist in a language, they were subject to forward and backward translation procedures. Well-validated measures of predictors and outcomes and items measuring COVID-19 related characteristics were selected after a consensus agreement among the members of this study.

## Predictors

**Country.** Respondents' countries were coded and entered as predictors.

**Socio-demographic status.** Participants responded to questions related to their socio-demographic characteristics including their age, gender, country of residence, marital status, employment status, educational level, whether they have children as well as their living situation.

**Lockdown variables.** Participants responded to questions related to lockdown including length of lockdown, whether they need to leave home for work, any change in their finances, whether they were able to obtain basic supplies, the amount of their living space confined in during the lockdown. They were also asked whether they, their partner, or a significant other was diagnosed with COVID-19.

**Social factors.** Social factors were measured using the Brief Assessment of Family Functioning Scale (BAFFS; [25]) and the Oslo Social Support Scale (OSSS; [26]). The BAFFS items are summed to produce a single score with higher scores indicating worse family functioning. The OSSS items are summed up and provide three levels types of social support: low (scored 3–8), moderate (scored 9–11) and high (scored 12–14).

**Psychological factors.** Psychological factors including mindfulness and psychological flexibility. Mindfulness was measured using the Cognitive Affective Mindfulness Scale (CAMS; [27]). The CAMS produces a single score with higher scores indicating better

mindfulness qualities. Psychological flexibility (e.g., hold one's thoughts lightly, be accepting of one's experiences, engage in what is important to them despite challenging situations) was measured using the Psyflex scale [28]. The Psyflex produces a single score with higher scores indicating better psychological flexibility qualities.

## Outcomes

**Stress.** Stress was measured using the Perceived Stress Scale (PSS; [29]). The PSS assesses an individual's appraisal of how stressful situations in their life are. Items ask about people's feelings and thoughts during the last month. A total score is produced, with higher scores indicating greater overall distress.

**Depression.** Depressive symptomatology was assessed using two items from the disengagement subscale of the Multidimensional State Boredom Scale (MSBS; [30]). These items assessed wanting to do pleasurable things but not finding anything appealing (i.e., boredom), as well as wasting time. Based on concepts of reinforcement deprivation (i.e., lack of access to or engagement with positive stimuli) that is known to contribute to depression, we added an item that measured how rewarding or pleasurable people found the activities that they were engaging in (i.e., reinforcement). Higher scores indicated higher depressive symptomatology.

**Positive affect/ negative affect.** The Positive And Negative Affect Scale (PANAS) was used to measure affect [31]. The original version of the questionnaire was used with five additional items: bored, confused, angry, frustrated and lonely. All items were scored on a 5-point Likert type scale, ranging from 1 = very little/not at all to 5 = extremely and summed up so that higher scores in the positive-related items indicating higher positive affect and higher scores in the negative-related items indicating higher negative affect. In order to capture additional dimensions of negative affect believed to be relevant to the COVID-19 lockdowns, we additionally added five items: bored, confused, angry, frustrated, lonely.

**Wellbeing.** Wellbeing was assessed using the Mental Health Continuum Short Form (MHC-SF; [32]); which assesses three aspects of wellbeing: emotional, psychological, and social. The MHC-SF produces a total score and scores for each of the three aspects of wellbeing. The MHC-SF can also be scored to produce categories of languishing (i.e., low levels of emotional, psychological, and social well-being), flourishing (i.e., high levels of emotional psychological and social well-being almost every day), and moderately mentally healthy (in between languishing and flourishing).

## Statistical analysis

The mean and standard deviation was calculated for dependent variables that follow the normal distribution while the median and interquartile range (IQR) were computed for non-normally distributed data. Bivariable association between an outcome variable and each predictor was investigated with ANOVA test for categorical predictor and univariable linear regression for numerical predictor. Linear mixed-effect model with random effect for country was performed to consider simultaneously several predictors in the same model and to account for the variation in outcome variable between countries. Four separate linear mixed-effect models were used for each outcome variable, one for each set of socio-demographic, lockdown, social and psychosocial predictors and multicollinearity for each set of predictors was investigated with the variation inflation criterion (VIF). Standardized regression coefficients were computed as effect size indices to measure the strength of the association between predictor variables and outcome variables. The comparison between the country mean and overall mean for each outcome variable was estimated through a linear regression model with dependent variable the mean centering outcome and predictor the country. Cohen's *d* effect size of the

standardize difference between country mean and the overall mean was computed as a measure of the magnitude of the difference between the two means.

The whole sample was used in linear mixed-effect models while for the comparison of country mean to the overall mean was used the sample from countries with sample size  $\geq 100$ . The R packages *lme4* and *effect sizes* were used for fitting the linear mixed effect model and to compute the standardized regression coefficients of the linear mixed effect models [33]. Significance test and confidence intervals were calculated at a significance level of 0.05. The following cut-off values were used for the evaluation of the effect sizes: 'tiny'  $\leq 0.05$ , 'very small' from 0.05 to  $\leq 0.10$ , 'small' from 0.10 to  $\leq 0.20$ , 'medium' from 0.20 to  $\leq 0.30$ , 'large' from 0.30 to  $\leq 0.40$  and 'very large'  $> 0.40$  [34].

## Results

### Descriptive

Participants were  $n = 9,565$  people from 78 countries. See [supporting information](#) for a participation flowchart ([S1 Appendix](#)). The countries with the largest samples were: Latvia ( $n = 1285$ ), Italy ( $n = 962$ ), Cyprus ( $n = 957$ ), Turkey ( $n = 702$ ), Switzerland ( $n = 550$ ), Hong Kong ( $n = 516$ ), Colombia ( $n = 485$ ), Ireland ( $n = 414$ ), Austria ( $n = 368$ ), Romania ( $n = 339$ ), Portugal ( $n = 334$ ), France ( $n = 313$ ), Spain ( $n = 296$ ), Germany ( $n = 279$ ), Hungary ( $n = 273$ ), Greece ( $n = 270$ ), USA ( $n = 268$ ), Finland ( $n = 157$ ), Montenegro ( $n = 147$ ), Poland ( $n = 135$ ), United Kingdom ( $n = 100$ ), Slovenia ( $n = 77$ ), and Canada ( $n = 60$ ). The remaining countries are listed in the supporting information ([S1 Table](#)).

### Outcome variables

The means, standard deviations, and where appropriate percentage of participants within categories of the five outcome variables can be seen in [Table 1](#).

### Predictor variables

**Countries.** A full list of countries can be found in the supporting information ([S1 Table](#)).

**Socio-demographic status.** The mean age was 36.9 (13.3) years. A majority of participants were female (77.7%), approximately a fifth male (22.0%), and small minority identified as other (0.3%). More than half of the respondents were either in a relationship (25.7%) or married (36.1%), almost a third were single (30.8%), and the rest were either divorced (5%), widower (1.1%) or other (1.3%). Participants indicated that they lived: alone (14.6%), with both parents (20.8%), one parent (5.1%), with their own family including partner and children (54.1%), or with friends or roommates (5.5%). Less than half of respondents had children (40.8%). Approximately half of the participants were working full time (53.4%), almost a fifth were working part-time (17.5%), 23.2% were unemployed and a small minority were either on parental leave (2.2%) or retired (3.7%).

**COVID-19 lockdown variables.** At the time of responding, participants were in lockdown or self-isolation for a median of 5.0 (3.0 IQR) weeks. Most people indicated that they had not been infected with COVID-19 (88.0%), a small minority indicated they had been infected (1.4%) and the rest had symptoms but were unsure (10.6%). Similar patterns were seen with reported infection rates of partners (no: 92.2%, yes: 0.7%, unsure: 7.1%) and of people close to them (no: 86.0%; yes: 5.6%; unsure: 8.4%). With respect to leaving the house for work, almost half (47.7%) indicated that this never occurred, 7.7% indicated leaving only once, whereas an almost equal number indicated leaving a couple times per week (23.7%) or more than three times per week (21.0%). Nearly all participants indicated they were able to obtain all the basic supplies they needed (93.5%). Participants reported having a median inner living

**Table 1. Descriptive statistics for outcomes and predictors (continuous variables).**

	Mean (M) (SD)	%
<b>Outcomes</b>		
Stress	17.1 (7.5)	
Low		33.0%
Moderate		55.9%
High		11.1%
Depression Symptoms	6.6 (2.3)	
Lack of Reinforcement	2.1 (1.0)	25.8% <sup>a</sup>
Boredom	2.4 (1.1)	32.4% <sup>a</sup>
Waste Time	2.4 (1.1)	47.4% <sup>a</sup>
Positive Affect	30.0 (8.1)	
Negative Affect	19.6 (7.8)	
Wellbeing	41.0 (14.0)	
Languishing		10.1%
Moderately Mentally Healthy		50.0%
Flourishing		39.9%
<b>Predictors</b>		
Family Functioning	5.5 (1.9)	
Social Support	9.9 (2.1)	
Mindfulness	26.7 (3.2)	
Psychological Flexibility	21.8 (4.1)	

Note: Possible ranges of the scales were as follows: Stress (PSS; 0–40); Reinforcement (1–4); Boredom (MSBS; 1–4); Positive Affect (PANAS-P; 10–50); Negative Affect (PANAS-N, plus; 15–75); Wellbeing (MHC-SF; 0–70); Family Functioning (BAFFS; 3–12); Social Support (OSSS; 3–14); Mindfulness (CAMS-R; 0–40); Psychological Flexibility (Psyflex; 6–30)

<sup>a</sup> = % of participants who responded with “lot” or “extremely”.

<https://doi.org/10.1371/journal.pone.0244809.t001>

space of 90.0 square meters (80.0 IQR) and median outdoor space of 20.0 square meters (192.1 IQR). Finally, with respect to finances, more than half indicated that their financial situation remained about the same (57.9%), a minority indicated it improved (8.9%), and a third reported that their finances had gotten worse (33.3%).

**Social and psychological predictors.** Mean values of the other predictors (i.e., social predictors and psychological predictors) can be seen in [Table 1](#).

## Multivariate analyses

**Stress.** Results of multivariate analyses for the outcome of stress can be seen in [Table 2](#). The largest protective factor against stress was social support (high support vs low support (-3.35, 95%CI, -3.39 to -2.92), with a very large effect size). Positive predictors of stress with large effect sizes were being female (2.42, 95%CI, 2.07 to 2.77) and worsening of finances (2.32, 95%CI, 1.68 to 2.96), whereas psychological flexibility buffered this response (-0.65, 95%CI, -0.69 to -0.62). Higher education levels were also associated with lower levels of stress, with a large effect size (see [Table 2](#)). Moderate effect sizes for predictors associated with less stress were older age (-0.13, 95%CI, -0.14, -0.11) and mindfulness (-0.69, 95%CI, -0.74, -0.64). Moderate effect sizes of predictors associated with more stress were worse family functioning (0.98, 95%CI, 0.90, 1.06) and not being able to obtain all basic supplies (1.82 95%CI, 1.12, 2.52).

Differences in reported levels of stress across countries were largely negligible, with the exception of two countries that reported higher levels of stress (Hong Kong (2.85, 95%CI, 2.22,



Table 2. Predictors of stress.

	PSS Score	
	Coefficient (95% CI)	Effect Size (95% CI)
<b>Socio Demographic Predictors</b>		
Sex		
Male	Ref	Ref
Female	2.42 (2.07, 2.77)	0.34 (0.29, 0.39)
Age	-0.13 (-0.14, -0.11)	-0.23 (-0.26, -0.19)
Employment		
Working (full time)	Ref	Ref
Working (part time)	0.23 (-0.18, 0.64)	0.03 (-0.02, 0.09)
Unemployed	1.08 (0.66, 1.51)	0.15 (0.09, 0.21)
On parental leave	-0.05 (-1.03, 0.94)	-0.01 (-0.14, 0.13)
Retired	-0.39 (-1.25, 0.58)	-0.05 (-0.17, 0.07)
Education		
Primary School	Ref	Ref
High School	-2.55 (-4.18, -0.92)	-0.36 (-0.60, -0.13)
Some College/University	-2.11 (-3.74, -0.47)	-0.30 (-0.53, -0.07)
Graduated from College/University	-2.49 (-4.10, -0.87)	-0.36 (-0.59, -0.13)
Master/Postgraduate	-2.81 (-4.42, -1.19)	-0.40 (-0.63, -0.17)
Doctoral level	-2.50 (-4.17, -0.82)	-0.36 (-0.60, 0.17)
Other	-1.78 (-3.60, 0.03)	-0.26 (-0.51, 0.00)
Marital status		
Single	Ref	Ref
In a relationship/engaged	0.20 (-0.24, 0.64)	0.03 (-0.03, 0.09)
Married	0.11 (-0.46, 0.69)	0.02 (-0.06, 0.10)
Divorced	-0.74 (-1.53, 0.03)	-0.10 (-0.21, 0.00)
Widower	0.50 (-0.95, 1.95)	0.07 (-0.13, 0.27)
Other	0.30 (-0.99, 1.59)	0.04 (-0.14, 0.22)
Living situation		
Live alone	Ref	Ref
Live with parents	-0.18 (-0.35, 0.73)	-0.03 (-0.10, 0.05)
Live with one of parents	0.96 (0.29, 1.80)	0.13 (0.03, 0.24)
Live with my own family	-0.70 (-1.37, -0.26)	-0.10 (-0.17, -0.02)
Live with friends/roommates	-0.05 (-0.66, 0.81)	-0.01 (-0.11, 0.09)
Having children		
Yes	Ref	Ref
No	-0.72 (-1.18, -0.26)	-0.10 (-0.16, -0.04)
<b>Lockdown Predictors</b>		
Weeks in quarantine/self-isolation	0.05 (-0.00, 0.14)	0.00 (0.00, 0.05)
Leave for work		
No	Ref	Ref
Once only	0.14 (-0.51, 0.79)	0.02 (-0.07, 0.11)
A couple of times	-0.97 (-1.41, -0.53)	-0.13 (-0.19, -0.07)
More than 3 times per week	-1.03 (-1.50, -0.56)	-0.14 (-0.21, -0.08)
Changes in finance		
Have gotten better	Ref	Ref
Stayed about the same	0.54 (-0.06, 1.15)	0.07 (-0.01, 0.15)
Have gotten worse	2.32 (1.68, 2.96)	0.31 (0.23, 0.40)

(Continued)

Table 2. (Continued)

	PSS Score	
	Coefficient (95% CI)	Effect Size (95% CI)
Obtain all basic supplies		
Yes	Ref	Ref
No	1.82 (1.12, 2.52)	0.24 (0.15, 0.34)
Inner living space (m2)	-0.02 (-0.03, -0.00)	-0.03 (-0.06, -0.01)
Outer space (m2)	-0.00 (-0.00, 0.00)	-0.01 (-0.03, 0.01)
<b>Social Predictors</b>		
Family Functioning Score	0.98 (0.90, 1.06)	0.24 (0.22, 0.26)
Perceived Social support		
Low	Ref	Ref
Moderate	-2.09 (-2.45, -1.72)	-0.27 (-0.32, -0.23)
High	-3.35 (-3.79, -2.92)	-0.44 (-0.49, -0.38)
<b>Psychological Predictors</b>		
Mindfulness	-0.69 (-0.74, -0.64)	-0.29 (-0.31, -0.27)
Psychological Flexibility	-0.65 (-0.69, -0.62)	-0.36 (-0.38, -0.34)

Note: Effect size interpretation: tiny (0–0.05); very small (0.05–0.10); small (0.10–0.20); medium (0.20–0.30); large (0.30–0.40); very large (>.40); Color code: light green–dark green (tiny–very large positive effect); light red–dark red (tiny–very large negative effect)

<https://doi.org/10.1371/journal.pone.0244809.t002>

3.49) and Turkey (2.47, 95%CI, 1.93, 3.02)) and two that reported lower levels of stress (Portugal (-2.50, 95%CI, -3.29, -1.71) and Montenegro (-3.30, 95%CI, -4.49, -2.11)) than the average stress level across all countries. See [supporting information](#) for information on each country (S2–S6 Tables).

**Depression.** Results of multivariate analyses for the outcome of depression can be seen in [Table 3](#). The strongest predictor of depression was social support, such that high (-1.30, 95% CI, -1.44, -1.16) and medium levels (-0.73, 95%CI, -0.85, -0.62) of social support were protective against depression (relative to low levels) with a very large and large effect sizes, respectively. The only other large effect size was for psychological flexibility, which also served in a protective manner (-0.20, 95%CI, -0.22, -0.19). Moderate effect sizes of predictors associated with less depression symptoms were also observed for higher education levels (see [Table 3](#)). Moderate effect sizes of predictors associated with more depression were worse family functioning (0.29, 95%CI, 0.27, 0.32) and not being able to obtain all basic supplies (0.49, 95%CI, 0.27, 0.70).

The amount of depression symptoms reported on average within countries was similar for most countries with the exception of one country with lower reported levels than average with a large effect size (Austria (-0.71, 95%CI, -0.95, -0.47)) and one with higher levels than average with a large effect size (USA (0.85, 95%CI, 0.58, 1.13)). See [supporting information](#) for information on each country (S2–S6 Tables).

**Affect.** Results of multivariate analyses for the outcome of affect can be seen in [Table 4](#). With respect to positive affect, social support (high support vs low support (5.69, 95%CI, 5.23, 6.16) and psychological flexibility (0.77, 95%CI, 0.74, 0.81) were both predictors with very large effect sizes. Interestingly, those who left their house more than three times per week had higher levels of positive affect than those that did not leave their house for work (1.68, 95%CI, 1.18, 2.17), with a medium effect size. Higher education levels were associated with higher levels of positive affect with a medium to large effect size (see [Table 4](#), PANAS-Positive).

Table 3. Predictors of depression.

	MSBS Score	
	Coefficient (95% CI)	Effect Size (95% CI)
<b>Socio Demographic Predictors</b>		
Sex		
Male	Ref	Ref
Female	0.08 (-0.03, 0.19)	0.04 (-0.01, 0.09)
Age	-0.03 (-0.03, -0.02)	-0.17 (-0.20, -0.14)
Employment		
Working (full time)	Ref	Ref
Working (part time)	-0.05 (-0.17, 0.08)	-0.02 (-0.08, 0.04)
Unemployed	0.28 (0.14, 0.40)	0.12 (0.06, 0.18)
On parental leave	-0.04 (-0.34, 0.26)	-0.02 (-0.16, 0.12)
Retired	0.24 (-0.02, 0.51)	0.11 (-0.01, 0.23)
Education		
Primary School	Ref	Ref
High School	-0.20 (-0.70, 0.30)	-0.09 (-0.32, 0.14)
Some College/University	-0.15 (-0.65, 0.35)	-0.09 (-0.32, 0.14)
Graduated from College/University	-0.44 (-0.94, 0.05)	-0.20 (-0.43, 0.02)
Master/Postgraduate	-0.62 (-1.11, -0.12)	-0.28 (-0.51, -0.05)
Doctoral level	-0.67 (-1.18, -0.16)	-0.31 (-0.54, -0.07)
Other	-0.43 (-0.99, 0.12)	-0.20 (-0.45, 0.06)
Marital status		
Single	Ref	Ref
In a relationship/engaged	-0.13 (-0.27, 0.00)	-0.06 (-0.13, 0.00)
Married	-0.16 (-0.34, 0.01)	-0.08 (-0.16, 0.01)
Divorced	0.04 (-0.20, 0.28)	0.02 (-0.09, 0.13)
Widower	-0.03 (-0.48, 0.41)	-0.02 (-0.23, 0.19)
Other	-0.06 (-0.45, 0.33)	-0.03 (-0.21, 0.16)
Living situation		
Live alone	Ref	Ref
Live with parents	-0.00 (-0.17, 0.16)	-0.00 (-0.08, 0.08)
Live with one of parents	0.09 (-0.14, 0.32)	0.04 (-0.06, 0.15)
Live with my own family	-0.16 (-0.33, 0.01)	-0.07 (-0.15, 0.00)
Live with friends/roommates	-0.24 (-0.47, -0.02)	-0.11 (-0.22, -0.01)
Having Children		
Yes	Ref	Ref
No	0.02 (-0.12, 0.16)	0.00 (-0.05, 0.07)
<b>Lockdown Predictors</b>		
Weeks in quarantine/self-isolation	-0.00 (-0.02, 0.02)	-0.00 (-0.03, 0.02)
Leave for work		
No	Ref	Ref
Once only	-0.12 (-0.31, 0.08)	-0.05 (-0.14, 0.04)
A couple of times	-0.19 (-0.32, -0.06)	-0.09 (-0.15, -0.03)
More than 3 times per week	-0.32 (-0.46, -0.18)	-0.15 (-0.21, -0.08)
Changes in finance		
Have gotten better	Ref	Ref
Stayed about the same	0.12 (-0.06, 0.30)	0.05 (-0.03, 0.13)
Have gotten worse	0.31 (0.12, 0.51)	0.14 (0.05, 0.23)

(Continued)

**Table 3.** (Continued)

	MSBS Score	
	Coefficient (95% CI)	Effect Size (95% CI)
Obtain all basic supplies		
Yes	Ref	Ref
No	0.47 (0.26, 0.68)	0.21 (0.12, 0.30)
Inner living space (m2)	-0.00 (-0.00, 0.00)	-0.04 (-0.06, -0.01)
Outer space (m2)	0.00 (-0.00, 0.00)	0.00 (-0.02, 0.02)
<b>Social Predictors</b>		
Family Functioning Score	0.28 (0.26, 0.30)	0.23 (0.21, 0.25)
Perceived Social support		
Low	Ref	Ref
Moderate	-0.69 (-0.80, -0.58)	-0.31 (-0.35, -0.26)
High	-1.23 (-1.36, -1.10)	-0.54 (-0.60, -0.48)
<b>Psychological Predictors</b>		
Mindfulness	-0.13 (-0.14, -0.11)	-0.18 (-0.20, -0.16)
Psychological Flexibility	-0.19 (-0.21, -0.18)	-0.35 (-0.37, -0.33)

Note: Effect size interpretation: tiny (0–0.05); very small (0.05–0.10); small (0.10–0.20); medium (0.20–0.30); large (0.30–0.40); very large (>.40); Color code: light green–dark green (tiny–very large positive effect); light red–dark red (tiny–very large negative effect)

<https://doi.org/10.1371/journal.pone.0244809.t003>

The amount of positive affect reported on average within countries was similar for most countries with the exception of one country with lower reported levels than average with a large effect size (Finland (-2.96, 95%CI, -4.19, -1.73)) and one with higher reported levels than average with a large effect size (Portugal (2.96, 95%CI, 2.12, 3.80)). See [supporting information](#) for information on each country (S2–S6 Tables).

With respect to negative affect, social support (high support vs low support (-2.74, 95%CI, -3.2, -2.29) and psychological flexibility (-0.62, 95%CI, -0.66, -0.58) were again the strongest associated predictors, with large effects. Higher education levels were also associated with lower levels of negative affect, with a medium effect (see [Table 4](#), PANAS-Negative). Higher levels of negative affect were noted, with medium effect sizes, for the predictors: worsening of finances (1.75, 95%CI, 1.10, 2.40) and not being able to obtain all basic supplies (1.6, 95%CI, 0.89, 2.31).

The amount of negative affect reported on average within countries was similar for most countries with the exception of few countries with lower reported negative affect levels than average with a very large effect sizes (Switzerland (-4.96, 95%CI, -5.91, -4.01), Germany (-4.70, 95%CI, -6.03, -3.37) & Austria (-6.49, 95%CI, -7.65, -5.33)) and one with a large effect size (Montenegro (-3.56, 95%CI, -5.39, -1.73)). The average amount of negative affect was higher than average in two countries, with very large effects size (Turkey (5.75, 95%CI, 4.92, 6.59) & Finland (7.57, 95%CI, 5.80, 9.34)). See [supporting information](#) for information on each country (S2–S6 Tables).

**Wellbeing.** Results of multivariate analyses for the outcome of wellbeing can be seen in [Table 5](#). Once again, social support (high support vs low support (13.20, 95%CI, 12.39, 14.01)) and psychological flexibility (1.42, 95%CI, 1.34, 1.49) were the predictors with the largest effect sizes (very large) on wellbeing. Higher education levels were associated with higher levels of wellbeing with a medium to large effect sizes (see [Table 5](#)). Medium negative effect sizes were noted for family functioning (-1.98, 95%CI, -2.12, -1.83) and inability to obtain all basic

**Table 4. Predictors of affect.**

	PANAS-Positive		PANAS-Negative	
	Coefficient (95% CI)	Effect Size (95% CI)	Coefficient (95% CI)	Effect Size (95% CI)
Socio Demographic Predictors				
Sex				
Male	Ref	Ref	Ref	Ref
Female	-1.19 (-1.57, -0.81)	-0.15 (-0.19, -0.10)	1.22 (0.85, 1.56)	0.16 (0.11, 0.21)
Age	0.09 (0.07, 0.11)	0.15 (0.12, 0.19)	-0.08 (-0.10, -0.06)	-0.15 (-0.18, -0.11)
Employment				
Working (full time)	Ref	Ref	Ref	Ref
Working (part time)	-0.41 (-0.86, 0.03)	-0.05 (-0.11, 0.00)	0.17 (-0.26, 0.61)	0.02 (-0.03, 0.08)
Unemployed	-1.13 (-1.59, -0.66)	-0.14 (-0.20, -0.08)	0.72 (0.27, 1.17)	0.10 (0.04, 0.15)
On parental leave	-0.24 (-1.31, 0.84)	-0.03 (-0.16, 0.11)	0.68 (-0.36, 1.72)	0.09 (-0.05, 0.23)
Retired	-1.12 (-2.06, -0.18)	-0.14 (-0.26, -0.02)	-0.28 (-1.19, 0.63)	-0.04 (-0.16, 0.08)
Education				
Primary School	Ref	Ref	Ref	Ref
High School	1.83 (0.06, 3.61)	0.23 (0.01, 0.46)	-2.16 (-3.87, -0.44)	-0.29 (-0.51, -0.06)
Some College/University	1.66 (-0.12, 3.43)	0.21 (-0.02, 0.43)	-1.75 (-3.47, -0.04)	-0.24 (-0.46, -0.01)
Graduated from College/University	1.98 (0.22, 3.73)	0.25 (0.03, 0.47)	-2.21 (-3.90, -0.51)	-0.30 (-0.52, -0.07)
Master/Postgraduate	2.41 (0.65, 4.44)	0.30 (0.08, 0.53)	-2.53 (-4.23, -0.83)	-0.35 (-0.57, -0.12)
Doctoral level	2.62 (0.79, 4.44)	0.33 (0.10, 0.56)	-2.09 (-3.85, -0.34)	-0.28 (-0.51, -0.04)
Other	1.87 (-0.11, 3.84)	0.24 (-0.01, 0.48)	-1.61 (-3.52, 0.29)	-0.21 (-0.47, 0.04)
Marital status				
Single	Ref	Ref	Ref	Ref
In a relationship/engaged	0.43 (-0.05, 0.90)	0.05 (-0.01, 0.12)	-0.12 (-0.58, 0.35)	-0.02 (-0.08, 0.05)
Married	0.32 (-0.31, 0.95)	0.04 (-0.04, 0.12)	-0.43 (-1.04, 0.17)	-0.06 (-0.14, 0.02)
Divorced	0.37 (-0.48, 1.22)	0.05 (-0.06, 0.16)	-0.89 (-1.71, -0.07)	-0.12 (-0.23, -0.01)
Widower	-0.22 (-1.79, 1.36)	-0.03 (-0.23, 0.17)	-0.65 (-2.18, 0.86)	-0.09 (-0.29, 0.11)
Other	-0.18 (-1.58, 1.22)	-0.02 (-0.20, 0.16)	-0.70 (-2.04, 0.65)	-0.09 (-0.27, 0.09)
Living situation				
Live alone	Ref	Ref	Ref	Ref
Live with parents	0.39 (-0.21, 1.00)	0.05 (-0.03, 0.13)	-0.01 (-0.56, 0.60)	-0.00 (-0.08, 0.08)
Live with one of parents	-0.31 (-1.13, 0.51)	-0.04 (-0.14, 0.06)	0.49 (-0.29, 1.28)	0.07 (-0.04, 0.17)
Live with my own family	0.69 (0.09, 1.29)	0.09 (0.01, 0.16)	-0.49 (-1.06, 0.09)	-0.06 (-0.14, 0.01)
Live with friends/roommates	1.13 (0.33, 1.93)	0.14 (0.04, 0.25)	0.15 (-0.62, 0.91)	0.02 (-0.08, 0.12)
Having children				
Yes	Ref	Ref	Ref	Ref
No	-0.61 (-1.11, -0.11)	-0.07 (-0.14, -0.01)	-1.23 (-1.71, -0.75)	-0.16 (-0.22, -0.10)
Lockdown predictors				
Weeks in quarantine /self-isolation	0.03 (-0.03, 0.09)	0.01 (-0.01, 0.04)	0.06 (-0.01, 0.11)	0.02 (0.00, 0.05)
Leave for work				
No	Ref	Ref	Ref	Ref
Once only	0.75 (0.06, 1.44)	0.10 (0.01, 0.19)	-0.48 (-1.13, 0.18)	-0.06 (-0.15, 0.02)
A couple of times	1.21 (0.74, 1.67)	0.16 (0.10, 0.22)	-0.73 (-1.17, -0.29)	-0.10 (-0.16, -0.04)
More than 3 times per week	1.68 (1.18, 2.17)	0.22 (0.15, 0.28)	-0.84 (-1.31, -0.37)	-0.11 (-0.18, -0.05)
Changes in finance				
Have gotten better	Ref	Ref	Ref	Ref
Stayed about the same	-0.56 (-1.20, 7.95)	-0.07 (-0.15, 0.01)	0.33 (-0.28, 0.94)	0.05 (-0.04, 0.13)
Have gotten worse	-1.03 (-1.71, -0.36)	-0.13 (-0.21, -0.04)	1.75 (1.10, 2.40)	0.24 (0.15, 0.33)

(Continued)

Table 4. (Continued)

	PANAS-Positive		PANAS-Negative	
	Coefficient (95% CI)	Effect Size (95% CI)	Coefficient (95% CI)	Effect Size (95% CI)
Obtain all basic supplies				
Yes	Ref	Ref	Ref	Ref
No	-1.34 (-2.09, -0.60)	-0.17 (-0.26, -0.08)	1.60 (0.89, 2.31)	0.21 (0.12, 0.30)
Inner living space (m2)	0.00 (0.00, 0.00)	0.07 (0.04, 0.09)	-0.00 (-0.00, 0.00)	-0.00 (-0.03, 0.02)
Outer space (m2)	0.00 (-0.00, 0.00)	0.03 (-0.02, 0.03)	-0.00 (-0.00, 0.00)	-0.02 (-0.05, 0.00)
Social Predictors				
Family Functioning Score	-0.81 (-0.90, -0.73)	-0.19 (-0.21, -0.17)	0.87 (0.79, 0.96)	0.21 (0.19, 0.23)
Perceived Social support				
Low	Ref	Ref	Ref	Ref
Moderate	3.11 (2.73, 3.49)	0.39 (0.34, 0.44)	-1.92 (-2.290, -1.54)	-0.22 (-0.27, -0.18)
High	5.69 (5.23, 6.16)	0.71 (0.65, 0.77)	-2.74(-3.20, -2.29)	-0.32 (-0.37, -0.27)
Psychological Predictors				
Mindfulness	0.59 (0.54, 0.64)	0.23 (0.21, 0.25)	-0.50 (-0.55, -0.45)	-0.21 (-0.23, -0.18)
Psychological Flexibility	0.77 (0.74, 0.81)	0.39 (0.37, 0.41)	-0.62 (-0.66, -0.58)	-0.33 (-0.35, -0.31)

Note: Effect size interpretation: tiny (0–0.05); very small (0.05–0.10); small (0.10–0.20); medium (0.20–0.30); large (0.30–0.40); very large (>.40); Color code: light green–dark green (tiny–very large positive effect); light red–dark red (tiny–very large negative effect).

<https://doi.org/10.1371/journal.pone.0244809.t004>

supplies (-3.27, 95%CI, -4.67, -1.87). Two medium positive effect sizes were observed: mindfulness (0.95, 95%CI, 0.86–1.04) and living with friends/roommates ((3.04, 95%CI, 1.59, 4.48), relative to living alone).

The level of wellbeing reported on average within countries was similar for most countries with the exception of three countries with higher levels with large effect sizes (Austria (4.95, 95%CI, 3.55, 6.34), Finland (5.24, 95%CI, 3.10, 7.38), & Portugal (4.59, 95%CI, 3.12, 6.05)) and two countries with lower levels of wellbeing than average with large (Italy (-4.36, 95%CI, -11.06, 2.35)) and very large effect sizes (Hong Kong (-6.84, 95%CI, -8.02, -5.66)). See [supporting information](#) for information on each country (S2–S6 Tables).

## Discussion

The COVID-19 is the largest pandemic in modern history. This study assessed nearly 10,000 participants across many countries to examine the impact of the pandemic and resultant governmental lockdown measures on mental health. During the height of the lockdown, the pandemic was experienced as at least moderately stressful for most people, and 11% reported the highest levels of stress. Symptoms of depression were also high, including 25% of the sample indicating that the things they did were not reinforcing, 33% reporting high levels of boredom, and nearly 50% indicating they wasted a lot of time. Consistent with symptoms of stress and depression, 10% of participants were psychologically languishing. These results suggest that there is a subgroup of people who are especially suffering and that in about 50% of the respondents' levels of mental health was only moderate. Previous studies have found that along with low levels, even moderate levels of mental health (which consists of only moderate levels of emotional, psychological, and social well-being) are associated with increased subsequent disability, productivity loss, and healthcare use [35–37]. Not everyone was suffering, however, as evidenced by the nearly 40% of participants who reported levels of mental health consistent with flourishing. The present results, while serious, do not point to more severe reactions observed in previous samples of selective quarantined individuals or groups [2]. Perhaps the

Table 5. Predictors of wellbeing.

	MHC-SF-Total Score	
	Coefficient (95% CI)	Effect Size (95% CI)
<b>Socio Demographic Predictors</b>		
Sex		
Male	Ref	Ref
Female	0.21 (-0.47, 0.91)	0.02 (-0.03, 0.07)
Age	0.13 (0.10, 0.17)	0.13 (0.10, 0.16)
Employment		
Working (full time)	Ref	Ref
Working (part time)	0.08 (-0.74, 0.91)	0.00 (-0.05, 0.07)
Unemployed	-1.95 (-2.80, -1.10)	-0.14 (-0.20, -0.08)
On parental leave	1.00 (-0.91, 2.92)	0.07 (-0.07, 0.21)
Retired	-0.43 (-2.13, 1.26)	-0.03 (-0.15, 0.09)
Education		
Primary School	Ref	Ref
High School	1.94 (-1.15, 5.03)	0.15 (-0.09, 0.38)
Some College/University	1.86 (-1.24, 4.97)	0.14 (-0.09, 0.38)
Graduated from College/University	2.36 (-0.69, 5.41)	0.18 (-0.05, 0.41)
Master/Postgraduate	3.34 (0.27, 6.40)	0.25 (0.02, 0.49)
Doctoral level	4.13 (0.95, 7.31)	0.31 (0.07, 0.56)
Other	2.65 (-0.83, 6.12)	0.20 (-0.06, 0.47)
Marital status		
Single	Ref	Ref
In a relationship/engaged	1.27 (0.39, 2.15)	0.10 (0.03, 0.16)
Married	1.99 (0.84, 3.14)	0.15 (0.06, 0.24)
Divorced	1.96 (0.41, 3.51)	0.15 (0.03, 0.26)
Widower	1.87 (-0.96, 4.71)	0.14 (-0.07, 0.36)
Other	-1.46 (-4.15, 1.22)	-0.11 (-0.31, 0.09)
Living Situation		
Live alone	Ref	Ref
Live with parents	2.22 (1.12, 3.32)	0.17 (0.08, 0.25)
Live with one of parents	-0.07 (-1.56, 1.40)	-0.00 (-0.12, 0.11)
Live with my own family	2.06 (0.96, 3.15)	0.15 (0.07, 0.24)
Live with friends/roommates	3.04 (1.59, 4.48)	0.23 (0.12, 0.34)
Having Children		
Yes	Ref	Ref
No	-1.11 (-2.02, -0.19)	-0.08 (-0.14, -0.01)
<b>Lockdown Predictors</b>		
Weeks in quarantine/self-isolation	-0.04 (-0.15, 0.07)	-0.00 (-0.04, 0.02)
Leave for work		
No	Ref	Ref
Once only	0.74 (-0.54, 2.01)	0.05 (-0.04, 0.15)
A couple of times	1.79 (0.94, 2.64)	0.13 (0.07, 0.20)
More than 3 times per week	2.56 (1.65, 3.47)	0.19 (0.12, 0.26)
Changes in finance		
Have gotten better	Ref	Ref
Stayed about the same	-0.96 (-2.10, 0.19)	-0.07 (-0.15, 0.01)
Have gotten worse	-2.17 (-3.39, -0.95)	-0.16 (-0.25, -0.07)

(Continued)

Table 5. (Continued)

	MHC-SF-Total Score	
	Coefficient (95% CI)	Effect Size (95% CI)
Obtain all basic supplies		
Yes	Ref	Ref
No	-3.27 (-4.67, -1.87)	-0.24 (-0.34, -0.14)
Inner living space (m2)	0.01 (0.00, 0.01)	0.09 (0.07, 0.12)
Outer space (m2)	0.00 (0.00, 0.00)	0.03 (0.00, 0.05)
<b>Social Predictors</b>		
Family Functioning Score	-1.98 (-2.12, -1.83)	-0.26 (-0.28, -0.24)
Perceived Social support		
Low	Ref	Ref
Moderate	7.78 (7.13, 8.43)	0.56 (0.52, 0.61)
High	13.20 (12.39, 14.01)	0.95 (0.90, 1.01)
<b>Psychological Predictors</b>		
Mindfulness	0.95 (0.86, 1.04)	0.22 (0.20, 0.24)
Psychological Flexibility	1.42 (1.34, 1.49)	0.42 (0.40, 0.44)

Note: Effect size interpretation: tiny (0–0.05); very small (0.05–0.10); small (0.10–0.20); medium (0.20–0.30); large (0.30–0.40); very large (>.40); Color code: light green–dark green (tiny–very large positive effect); light red–dark red (tiny–very large negative effect)

<https://doi.org/10.1371/journal.pone.0244809.t005>

previously reported distress in these groups is prevented when an entire country or world is in lockdown so that the feeling emerges that “everyone is in it together”.

Importantly, a handful of predictors emerged that consistently predicted all outcomes: Social support, education level, finances, access to basic needs, and the ability to respond psychologically flexible. The consistency of results examining predictors is noteworthy, both in terms of the consistently strong predictors (e.g., social support, education, psychological flexibility, as well as loss of income and lack of access to necessities) and in terms of the other predictors that were either not predictive or only weakly so. All predictors were chosen based on theoretical ties to the outcomes, previous findings, and studies on quarantines [2].

A novel finding was that people who left their house three or more times per week reported more positive affect than those that left their house less often. It is possible that these people experienced more variation, which contributed to positive affect. It is also possible they experienced a greater sense of normality. Future studies are encouraged to further investigate possible mechanisms through which this result unfolds.

Overall, these patterns did not differ substantially between countries. Although some differences did emerge, they were mostly inconsistent across outcomes. Three countries fared worse on two outcomes each: Hong Kong (stress & wellbeing); Turkey (stress & negative affect); and Finland (lower positive affect and higher negative affect)—though participants in Finland also reported higher levels of wellbeing than average. Two countries had more favorable outcomes than the average levels across all countries: Portugal (lower stress and higher wellbeing) and Austria (lower depression and higher wellbeing). The differences observed are likely due to a combination of chance, sampling, nation specific responses to the COVID-19 pandemic, cultural differences, and other factors playing out in the countries (e.g., political unrest [38]). If replicated, future studies are encouraged to examine possible mechanisms of these outcomes.

This study provides valuable insights on several levels. First, it documents the mental health outcomes across a broad sample during the COVID-19 global pandemic. Second, it informs about the conditions and resilience factors (social support, education, and psychological



flexibility) and risk factors (loss of income and inability to get basic supplies) that affect mental health outcomes. Third, these factors can be used in future public health responses are being made, including those that require large scale lockdowns or quarantines. That is, public health officials should direct resources to identifying and supporting people with poor social support, income loss, and potentially lower levels of education and provide a strategy to mitigate special risks in these subpopulations. The importance of social support needs to be made clear to the public and to the degree possible mechanisms that can contribute to social support should be supported. Further, psychological flexibility is a trainable set of skills that has repeatedly been shown to ameliorate suffering [22, 39]; and can be widely distributed with modern technological intervention tools such as digital, internet, or virtual means [40]. We do not claim, however, that psychological flexibility is the only factor that can be used for interventions. Instead, it is a recognized transdiagnostic factor assessed in this study and one that is feasible to be targeted and modified by interventions and prevention [41–43].

This study is limited by several important factors. First, the results are based on cross sectional analysis and correlations. As such, causation cannot be inferred and any delayed impact of the pandemic and lockdown on peoples' mental health was not captured. Second, all results of this survey were obtained via self-report questionnaires, which can be subject to retrospective response bias. Third, although the sample was large and based on varied recruitment sources, it was not representative of the population and undersampled people who suffered most from the pandemic (i.e., front line health care professionals, people in intensive care, etc.) or people without internet access, etc. Finally, the country-specific incidence rates and lockdown measures differed across countries. These were not assessed, but future studies are encouraged to investigate how such factors impact mental health outcomes.

These limitations notwithstanding, based on nearly 10,000 international participants, this study found that approximately 10% of the population was languishing during or shortly after the lockdown period. These finding have implications for public health initiatives. First, officials are urged to attend to, find, and target people who have little social support and/ or whose finances have worsened as a result of the measures. Second, public health interventions are further urged to target psychological processes such as psychological flexibility in general to potentially help buffer other risk factors for mental health. Likewise, availability of social support and information about where to get support and remain connected are needed. These recommendations should become part of public health initiatives designed to promote mental health in general, and should equally be considered when lockdowns or physical distancing are prescribed during a pandemic.

## Supporting information

**S1 Table. List of all countries included in the data set.**

(PDF)

**S2 Table. Geodemographic predictors for Perceived Stress Scale.**

(PDF)

**S3 Table. Geodemographic predictors for MSBS–depression.**

(PDF)

**S4 Table. Geodemographic predictors for PANAS positive.**

(PDF)

**S5 Table. Geodemographic predictors for PANAS negative.**

(PDF)

**S6 Table. Geodemographic predictors for MHCSF—mental health continuum.**  
(PDF)

**S1 Appendix. Participation flowchart.**  
(PDF)

## Acknowledgments

We wish to thank the following people for their work in helping to implement the study: Spyros Demosthenous, Christiana Karashali, Diamanto Rovania (University of Cyprus); Maria Antoniadou (European University of Cyprus); Ioanna Menoikou (Cyprus University of Technology); Elias Ioannou (University of Nicosia); Sonja Borner, Victoria Firsching-Block, Alexander Fenn (University of Basel); Cristine Šneidere, Ingrida Trups-Kalne, Lolita Vansovica, Sandra Feldmane, (Riga Stradiņš University); David Nilsson (Lund University); Miguel A. Segura-Vargas (Fundación Universitaria Konrad Lorenz); Claudia Lenuța Rus, Catalina Otoi, Cristina Vajaeon (Babes-Bolyai University). We further wish to thank Fabio Coviello and Sonja Borner (University of Basel) for their help in preparing the manuscript.

## Author Contributions

**Conceptualization:** Andrew T. Gloster, Maria B. Garcia-Martin, Angelos P. Kassianos, Maria Karekla.

**Data curation:** Demetris Lamnisos.

**Formal analysis:** Demetris Lamnisos.

**Funding acquisition:** Andrew T. Gloster.

**Investigation:** Jelena Lubenko, Giovambattista Presti, Valeria Squatrito, Marios Constantinou, Christiana Nicolaou, Savvas Papacostas, Gökçen Aydın, Yuen Yu Chong, Wai Tong Chien, Ho Yu Cheng, Francisco J. Ruiz, Maria B. Garcia-Martin, Diana P. Obando-Posada, Miguel A. Segura-Vargas, Vasilis S. Vasiliou, Louise McHugh, Stefan Höfer, Adriana Baban, David Dias Neto, Ana Nunes da Silva, Jean-Louis Monestès, Javier Alvarez-Galvez, Marisa Paez-Blarrina, Francisco Montesinos, Sonsoles Valdivia-Salas, Dorottya Ori, Bartosz Kleszcz, Raimo Lappalainen, Iva Ivanović, David Gosar, Frederick Dionne, Rhonda M. Merwin.

**Methodology:** Andrew T. Gloster, Angelos P. Kassianos, Maria Karekla.

**Project administration:** Andrew T. Gloster, Angelos P. Kassianos, Maria Karekla.

**Resources:** Jelena Lubenko, Giovambattista Presti, Valeria Squatrito, Marios Constantinou, Christiana Nicolaou, Savvas Papacostas, Gökçen Aydın, Yuen Yu Chong, Wai Tong Chien, Ho Yu Cheng, Francisco J. Ruiz, Maria B. Garcia-Martin, Diana P. Obando-Posada, Miguel A. Segura-Vargas, Vasilis S. Vasiliou, Louise McHugh, Stefan Höfer, Adriana Baban, David Dias Neto, Ana Nunes da Silva, Jean-Louis Monestès, Javier Alvarez-Galvez, Marisa Paez-Blarrina, Francisco Montesinos, Sonsoles Valdivia-Salas, Dorottya Ori, Bartosz Kleszcz, Raimo Lappalainen, Iva Ivanović, David Gosar, Frederick Dionne, Rhonda M. Merwin.

**Supervision:** Andrew T. Gloster, Angelos P. Kassianos, Maria Karekla.

**Writing – original draft:** Andrew T. Gloster, Demetris Lamnisos, Angelos P. Kassianos, Maria Karekla.

**Writing – review & editing:** Andrew T. Gloster, Demetris Lamnisos, Jelena Lubenko, Giovambattista Presti, Valeria Squatrito, Marios Constantinou, Christiana Nicolaou, Savvas

Papacostas, Gökçen Aydın, Yuen Yu Chong, Wai Tong Chien, Ho Yu Cheng, Francisco J. Ruiz, Maria B. Garcia-Martin, Diana P. Obando-Posada, Miguel A. Segura-Vargas, Vasilis S. Vasiliou, Louise McHugh, Stefan Höfer, Adriana Baban, David Dias Neto, Ana Nunes da Silva, Jean-Louis Monestès, Javier Alvarez-Galvez, Marisa Paez-Blarrina, Francisco Montecosinos, Sonsoles Valdivia-Salas, Dorottya Ori, Bartosz Kleszcz, Raimo Lappalainen, Iva Ivanović, David Gosar, Frederick Dionne, Rhonda M. Merwin, Angelos P. Kassianos, Maria Karekla.

## References

1. Presti G, McHugh L, Gloster A, Karekla M, Hayes SC. The dynamics of fear at the time of COVID-19: A contextual behavioral science perspective. *Clin Neuropsychiatry*. 2020; 17(2):65–71.
2. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. Vol. 395, *The Lancet*. Lancet Publishing Group; 2020. p. 912–20.
3. Kristensen P, Weisaeth L, Heir T. Bereavement and mental health after sudden and violent losses: A Review. Vol. 75, *Psychiatry*. Guilford Publications Inc; 2012. p. 76–97. <https://doi.org/10.1521/psyc.2012.75.1.76> PMID: 22397543
4. Lawrence BJ, Jayakody DMP, Bennett RJ, Eikelboom RH, Gasson N, Friedland PL. Hearing loss and depression in older adults: A systematic review and meta-analysis. *Gerontologist*. 2020. <https://doi.org/10.1093/geront/gnz009> PMID: 30835787
5. van Eersel JHW, Taris TW, Boelen PA. Reciprocal relations between symptoms of complicated grief, depression, and anxiety following job loss: A cross-lagged analysis. *Clin Psychol*. 2020 Mar;
6. Alfano CA, Bower JL, Cowie J, Lau S, Simpson RJ. Long-duration space exploration and emotional health: Recommendations for conceptualizing and evaluating risk. Vol. 142, *Acta Astronautica*. Elsevier Ltd; 2018. p. 289–99.
7. Berry JW, Hou F. Immigrant acculturation and wellbeing in Canada. *Can Psychol*. 2016 Nov; 57(4):254–64.
8. Shankar A, Rafnsson SB, Steptoe A. Longitudinal associations between social connections and subjective wellbeing in the English Longitudinal Study of Ageing. *Psychol Health*. 2015 Jun; 30(6):686–98. <https://doi.org/10.1080/08870446.2014.979823> PMID: 25350585
9. Kanter JW, Busch AM, Weeks CE, Landes SJ. The nature of clinical depression: Symptoms, syndromes, and behavior analysis. Vol. 31, *Behavior Analyst*. Association for Behavior Analysis; 2008. p. 1–21.
10. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and Social Isolation as Risk Factors for Mortality: A Meta-Analytic Review. *Perspect Psychol Sci*. 2015 Mar; 10(2):227–37. <https://doi.org/10.1177/1745691614568352> PMID: 25910392
11. Ryan RM, Bernstein JH, Brown KW. Weekends, work, and well-being: Psychological need satisfactions and day of the week effects on mood, vitality, and physical symptoms. *J Soc Clin Psychol*. 2010 Jan; 29(1):95–122.
12. Koenig HG. Research on religion, spirituality, and mental health: A review. Vol. 54, *Canadian Journal of Psychiatry*. Canadian Psychiatric Association; 2009. p. 283–91. <https://doi.org/10.1177/070674370905400502> PMID: 19497160
13. Paul KI, Moser K. Unemployment impairs mental health: Meta-analyses. *J Vocat Behav*. 2009 Jun; 74(3):264–82.
14. Lund C, Myer L, Stein DJ, Williams DR, Flisher AJ. Mental illness and lost income among adult South Africans. *Soc Psychiatry Psychiatr Epidemiol*. 2013 May; 48(5):845–51. <https://doi.org/10.1007/s00127-012-0587-5> PMID: 23007296
15. Phillips DR, Siu OL, Yeh AGO, Cheng KHC. Informal social support and older persons' psychological well-being in Hong Kong. *J Cross Cult Gerontol*. 2008 Mar; 23(1):39–55. <https://doi.org/10.1007/s10823-007-9056-0> PMID: 18228121
16. Golden J, Conroy RM, Bruce I, Denihan A, Greene E, Kirby M, et al. Loneliness, social support networks, mood and wellbeing in community-dwelling elderly. *Int J Geriatr Psychiatry*. 2009 Jul; 24(7):694–700. <https://doi.org/10.1002/gps.2181> PMID: 19274642
17. Gaymu J, Springer S. Living conditions and life satisfaction of older Europeans living alone: A gender and cross-country analysis. *Ageing Soc*. 2010 Oct; 30(7):1153–75.

18. Panayiotou G, Karekla M. Perceived social support helps, but does not buffer the negative impact of anxiety disorders on quality of life and perceived stress. *Soc Psychiatry Psychiatr Epidemiol*. 2013 Feb; 48(2):283–94. <https://doi.org/10.1007/s00127-012-0533-6> PMID: 22711064
19. Phillips TM. The Influence of Family Structure Vs. Family Climate on Adolescent Well-Being. *Child Adolesc Soc Work J*. 2012 Apr; 29(2):103–10.
20. Schrodtt P, Witt PL, Messersmith AS. A Meta-Analytical Review of Family Communication Patterns and their Associations with Information Processing, Behavioral, and Psychosocial Outcomes. *Commun Monogr*. 2008 Sep; 75(3):248–69.
21. Chiesa A, Serretti A. Mindfulness-based stress reduction for stress management in healthy people: A review and meta-analysis. *J Altern Complement Med*. 2009 May; 15(5):593–600. <https://doi.org/10.1089/acm.2008.0495> PMID: 19432513
22. Gloster AT, Meyer AH, Lieb R. Psychological flexibility as a malleable public health target: Evidence from a representative sample. *J Context Behav Sci*. 2017; 6(2):166–71.
23. Gu J, Strauss C, Bond R, Cavanagh K. How do mindfulness-based cognitive therapy and mindfulness-based stress reduction improve mental health and wellbeing? A systematic review and meta-analysis of mediation studies. Vol. 37, *Clinical Psychology Review*. Elsevier Inc.; 2015. p. 1–12. <https://doi.org/10.1016/j.cpr.2015.01.006> PMID: 25689576
24. Panayiotou G, Karekla M, Mete I. Dispositional coping in individuals with anxiety disorder symptomatology: Avoidance predicts distress. *J Context Behav Sci*. 2014 Oct; 3(4):314–21.
25. Mansfield AK, Keitner GI, Sheeran T. The Brief Assessment of Family Functioning Scale (BAFFS): a three-item version of the General Functioning Scale of the Family Assessment Device. *Psychother Res*. 2019 Sep; 29(6):824–31. <https://doi.org/10.1080/10503307.2017.1422213> PMID: 29351729
26. Dalgard O. Community health profile as tool for psychiatric prevention. *Promot Ment Heal*. 1996; 5:681–95.
27. Feldman G, Hayes A, Kumar S, Greeson J, Laurenceau JP. Mindfulness and emotion regulation: The development and initial validation of the Cognitive and Affective Mindfulness Scale-Revised (CAMS-R). *J Psychopathol Behav Assess*. 2007 Sep; 29(3):177–90.
28. Gloster AT, Block VJ, Klotsche J, Villanueva J, Rinner MTB, Benoy C, et al. Psy-Flex: A Contextually Sensitive State Measure of Psychological Flexibility. In review.
29. Cohen S. Perceived stress in a probability sample of the United States. In: *The social psychology of health*. Thousand Oaks, CA, US: Sage Publications, Inc; 1988. p. 31–67. (The Claremont Symposium on Applied Social Psychology.)
30. Fahlman SA, Mercer-Lynn KB, Flora DB, Eastwood JD. Development and validation of the multidimensional state boredom scale. *Assessment*. 2013 Feb; 20(1):68–85. <https://doi.org/10.1177/1073191111421303> PMID: 21903683
31. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol*. 1988; 54(6):1063. <https://doi.org/10.1037//0022-3514.54.6.1063> PMID: 3397865
32. Keyes CLM, Wissing M, Potgieter JP, Temane M, Kruger A, van Rooy S. Evaluation of the mental health continuum-short form (MHC-SF) in Setswana-speaking South Africans. *Clin Psychol Psychother*. 2008 May; 15(3):181–92. <https://doi.org/10.1002/cpp.572> PMID: 19115439
33. Bates D, Mächler M, Bolker BM, Walker SC. Fitting linear mixed-effects models using lme4. *J Stat Softw*. 2015 Oct; 67(1):1–48.
34. Funder DC, Ozer DJ. Evaluating Effect Size in Psychological Research: Sense and Nonsense. *Adv Methods Pract Psychol Sci*. 2019 Jun; 2(2):156–68.
35. Keyes CLM. The mental health continuum: From languishing to flourishing in life. *J Health Soc Behav*. 2002; 43(2):207–22. PMID: 12096700
36. Keyes CLM, Grzywacz JG. Health as a Complete State: The Added Value in Work Performance and Healthcare Costs. *J Occup Environ Med*. 2005 May; 47(5):523–32. <https://doi.org/10.1097/01.jom.0000161737.21198.3a> PMID: 15891532
37. Keyes CLM. Promoting and Protecting Mental Health as Flourishing: A Complementary Strategy for Improving National Mental Health. *Am Psychol*. 2007 Feb; 62(2):95–108. <https://doi.org/10.1037/0003-066X.62.2.95> PMID: 17324035
38. Ni MY, Yao XI, Leung KSM, Yau C, Leung CMC, Lun P, et al. Depression and post-traumatic stress during major social unrest in Hong Kong: a 10-year prospective cohort study. *Lancet*. 2020 Jan; 395(10220):273–84. [https://doi.org/10.1016/S0140-6736\(19\)33160-5](https://doi.org/10.1016/S0140-6736(19)33160-5) PMID: 31928765
39. Gloster AT, Klotsche J, Ciarrochi J, Eifert G, Sonntag R, Wittchen HU, et al. Increasing valued behaviors precedes reduction in suffering: Findings from a randomized controlled trial using ACT. *Behav Res Ther*. 2017 Apr; 91:64–71. <https://doi.org/10.1016/j.brat.2017.01.013> PMID: 28160720

40. Karekla M, Stella, Savvides N, Gloster A. An Avatar-Led Intervention Promotes Smoking Cessation in Young Adults: A Pilot Randomized Clinical Trial. *Ann Behav Med.* 2020; <https://doi.org/10.1093/abm/kaaa013> PMID: 32383736
41. Thompson E, Destree L, Albertella L, Fontenelle L. Internet-Based Acceptance and Commitment Therapy: A Transdiagnostic Systematic Review and Meta-Analysis for Mental Health Outcomes. *Behav Ther.* 2020 Jul;
42. Levin ME, MacLane C, Daflos S, Seeley JR, Hayes SC, Biglan A, et al. Examining psychological inflexibility as a transdiagnostic process across psychological disorders. *J Context Behav Sci.* 2013 Dec; 3 (3):155–63.
43. Dindo L, Van Liew JR, Arch JJ. Acceptance and Commitment Therapy: A Transdiagnostic Behavioral Intervention for Mental Health and Medical Conditions. Vol. 14, *Neurotherapeutics*. Springer New York LLC; 2017. p. 546–53. <https://doi.org/10.1007/s13311-017-0521-3> PMID: 28271287